arXiv:1304.2893v1 [math.DS] 10 Apr 2013

On embeddability of automorphisms into measurable flows from the point of view of self-joining properties

Joanna Kułaga-Przymus

April 11, 2013

Abstract

We compare self-joining- and embedability properties. In particular, we prove that a measure preserving flow $(T_t)_{t\in\mathbb{R}}$ with T_1 ergodic is 2-fold quasi-simple (2-fold distally simple) if and only if T_1 is 2-fold quasi-simple (2-fold distally simple). We also show that the Furstenberg-Zimmer decomposition for a flow $(T_t)_{t\in\mathbb{R}}$ with T_1 ergodic with respect to any flow factor is the same for $(T_t)_{t\in\mathbb{R}}$ and for T_1. We give an example of a 2-fold quasi-simple flow disjoint from simple flows and whose time-one map is simple. We describe two classes of flows (flows with minimal self-joining property and flows with the so-called Ratner property) whose time-one maps have unique embeddings into measurable flows. We also give an example of a 2-fold simple flow whose time-one map has more than one embedding.

Contents

1 Introduction **2**
1.1 Embeddability . . . 2
1.2 Joining properties . . . 3
1.3 Motivation and results . . . 5
1.3.1 Between simplicity and JP: 2-QS and 2-DS . . . 5
1.3.2 Mixing properties . . . 6
1.3.3 (Non-)uniqueness of the embedding . . . 7
1.3.4 Counter-examples . . . 8

2 Preliminaries **9**
2.1 Topology, invariant measures, ergodic decomposition . . . 10
2.2 Factors and extensions I . . . 11
2.3 Joinings I . . . 11
2.3.1 Basic definitions and properties . . . 11
2.4 Factors and extensions II . . . 13
2.4.1 Relatively weakly mixing extensions . . . 13
2.4.2 Relatively isometric and relatively distal extensions . . . 13
2.5 Joinings II . . . 14

Research supported by Narodowe Centrum Nauki grant DEC-2011/03/B/ST1/00407.

2.5.1 Self-joining properties 14
2.6 Distal and isometric extensions – tools 16
2.6.1 Separating sieves 16
2.6.2 T-compactness 17

3 Self-joining properties of $\mathcal{T}$ and T_1 – part I **17**
3.1 Simplicity 17
3.2 Joining primeness property 19
3.3 Isometric, distal and relatively weakly mixing extensions 22
3.3.1 Isometric extensions 22
3.3.2 Distal extensions and Furstenberg-Zimmer decomposition 27
3.4 Quasi- and distal simplicity 32
3.4.1 Quasi-simplicity – general case 32
3.4.2 Quasi-simplicity – special case 33
3.4.3 Distal simplicity 33

4 (Non)-uniqueness of the embedding **34**
4.1 Uniqueness 34
4.2 Non-uniqueness 37

5 Self-joining properties of $\mathcal{T}$ and T_1 – part II **39**
5.1 Easy counter-example 40
5.2 Advanced counter-example 42
5.2.1 (C, F)-constructions 42
5.2.2 Construction of a counter-example 45
5.2.3 Weak mixing 49
5.2.4 Self-joining properties 53

6 Appendix – uniform distribution **58**
6.1 Necessary tools 58
6.2 Technical results 60

1 Introduction

1.1 Embeddability

The problem of embeddability of automorphisms into a measurable flow[1] has already been studied in ergodic theory for more than 60 years. Even today, the basic problem – a necessary and sufficient condition for embeddability – remains the most interesting open question in this area. Historically, the first articles on this subject dealt with a simpler problem, namely the one of the existence of roots of ergodic automorphisms. In [31] Halmos gave a necessary condition for the existence of a square root (the absence of -1 in the spectrum). In the discrete spectrum case this is also sufficient. His result was later generalized in various ways – several authors considered roots of other orders both in the discrete [42, 5, 39, 61] and quasi-discrete spectrum [30, 50, 51, 52] case. These results include necessary and sufficient conditions for the existence of roots and

[1] By embeddability we will always mean embeddability into a measurable flow.

for the existence of embedding into a flow in the discrete and quasi-discrete spectrum case.[2] In particular, Lamperti [42] gave an example of a discrete spectrum automorphism with roots of all orders, which is however not embeddable into a flow.

The problem turned out to be much more challenging in the weakly mixing case. Already in 1956, Halmos asked [32] ([33]) about the existence of square roots of weakly mixing automorphisms, the existence of roots for Bernoulli shifts and the embeddability of Bernoulli shifts into flows. The answers to his questions came relatively fast. Chacon constructed a non-mixing automorphism with continuous spectrum having no square root [7] and a bit later one having no roots [8]. Since the centralizer of an automorphism embeddable into a flow contains all time-t automorphisms of this flow, a necessary condition for embeddability is that the centralizer of the considered automorphism is uncountable. Del Junco [15], continuing the study of the Chacon's type constructions, showed, in particular, that the centralizer of the classical Chacon's automorphism [9] consists only of its powers, whence it is too small for the automorphism even to admit roots. Further results were provided by Ornstein – he answered positively the remaining two questions of Halmos on the Bernoulli embedding [55, 57]. While all previous constructions were not mixing, and Bernoulli shifts have strong mixing properties, it seemed natural that suficiently strong mixing properties could imply embeddability (in 1966 this problem was still open [5]). This turned out to be false. Ornstein consecutively constructed a mixing automorphism [58] and a K-automorphism [60] without square roots.

After that there seems to have been no substantial progress until 2000 when King [37] showed that a generic automorphism admits roots of all orders. Three years later de la Rue and de Sam Lazaro proved even more – that a generic automorphism can be embedded into a flow [14]. This result was later generalized (in various ways) to the case of $\mathbb{Z}^d$-actions [73, 74, 69]. In 2004 Stepin and Eremenko [71] showed that typically there are uncountably many pairwise spectrally non-isomorphic embeddings, thus strengthening the result from [14].

1.2 Joining properties

We will deal with the problem of embeddability from the point of view of joining theory. Joinings were introduced to ergodic theory by Furstenberg [26] and have been a very fruitful tool for studying the dynamical systems from the very beginning. A joining of two (or more) systems (in our case automorphisms or flows) is a way these systems are seen as factors of a larger system, i.e. it describes the relation between the sub-σ-algebras corresponding to these two (or more) systems "placed" together in a larger one. More formally, a joining is a measure invariant under the product action whose marginals are the invariant measures of the original systems. We deal with self-joinings, i.e. joinings of two (or more) copies of a fixed system. Depending on the "number of ways" a system can be joined with itself in an ergodic way, one defines several classes of dynamical systems. The paper which initiated a systematic study of the self-joining properties was published by Rudolph [67], who introduced the notion of *minimal self-joinings* (MSJ). It is the most "restrictive" class of systems from

[2] Hahn and Parry [30] showed that automorphisms with quasi-discrete, non-discrete spectrum are not embeddable into flows.

this point of view. The only ergodic self-joinings of systems in this class (other than the product measure) are the so-called off-diagonal self-joinings, which are concentrated on graphs of automorphisms "contributing" to the action under consideration. Since for a typical automorphism the centralizer is uncountable (clearly, this is not the case for automorphisms with the MSJ property), it is natural to relax the assumptions and consider the so-called *2-fold simple* systems (Veech [75], del Junco, Rudolph [19]).[3] Here as ergodic self-joinings one also admits off-diagonal measures concentrated on the graphs of automorphisms from the centralizer. A yet larger class is the class of *2-fold quasi-simple (2-QS)* systems (Ryzhikov, Thouvenot [68]) and the class of *2-fold distally simple (2-DS)* systems (del Junco, Lemańczyk [18]). In the first of these classes as ergodic self-joinings one additionally admits joinings which are *relatively isometric* over the marginal factors and in the latter one – also the *relatively distal* ones (for the definitions see Section 2.4.2).[4] A particular case of the 2-QS systems are systems whose ergodic self-joinings other than the product measure are finite extensions of the marginal factors. We will denote them by "$n : 1$" for $n \geq 1$. For $n = 1$ this notion is equivalent to simplicity. The largest of the classes which will be of our interest are systems with *joining primeness property (JP)* (Lemańczyk, Parreau, Roy [45]). Contrary to the previous classes, systems with the JP property were originally no longer defined in terms of self-joinings – here some restriction is imposed on the type of ergodic joinings with Cartesian products of weakly mixing systems.[5]

One more property we will consider is the so-called *R-property*, which was first observed by Ratner in [66] for horocycle flows. It describes the behavior of orbits of distinct points which are close to each other. Roughly speaking, sufficiently long pieces of their orbits are close to each other up to a time shift. An important ingredient of the definition of the R-property is the restriction on how large this time shift can be. Precise conditions (more restrictive in [66], see also [24] and [25]) form a variety of combinatorial properties, all yielding the same dynamical consequences in case of weakly mixing flow: each of them implies that the flow is "$n : 1$" for some $n \in \mathbb{N}$.[6] In fact it implies more: ergodic joinings with other flows, other than the product measure, are also finite extensions of the coordinate factor of the "additional" flow [66] (see also [72]). Apart from the horocycle flows, several classes of flows satisfying the R-property are known. They are defined in terms of special flows: examples from [24, 35] are completely different from horocycle flows from the dynamical point of view as they are disjoint from all mixing flows; see also [25] for other examples.

The relations between the above-listed classes of systems can be summarized as follows:

[3] We will be mostly interested in 2-fold joinings. The self-joining properties of higher order are defined using the so-called PID property, see Section 2.5.1. Danilenko showed [11] that $\mathcal{T}$ has the PID property if and only if T_1 has this property.

[4] Recall that each 2-fold DS-system is either weakly mixing or it is distal.

[5] We show that the JP property is also an intrinsic one – it suffices to deal with products of systems which are factors of ergodic infinite self-joinings of the system under consideration.

[6] Another version of the R-property can be found in [76] – here however the notion was investigated from other point of view.



1.3 Motivation and results

1.3.1 Between simplicity and JP: 2-QS and 2-DS

A starting point for our investigations is a result of del Junco and Rudolph [20] who showed that whenever an ergodic flow is 2-fold simple then its time-one map is also 2-fold simple. On the other hand, it is not difficult to prove that given a flow $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ with T_1 ergodic, $\mathcal{T}$ has the JP property whenever T_1 has the JP property (see Section 3.2). Analogous results hold also for automorphisms and their powers.

The 2-QS and 2-DS properties are "intermediate" between simplicity and the JP property. Hence it is reasonable to expect that the 2-QS and 2-DS property will share either the features of the simple systems or the features of the JP systems, or both. Moreover, it is quite easy to see that if T_1 is 2-fold simple then $\mathcal{T}$ is 2-QS. This makes it even less surprising that indeed either both $\mathcal{T}$ and T_1 are 2-QS (2-DS) or none of them is:

Theorem 1.1. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be an ergodic flow. The following conditions are equivalent:*

(i) $\mathcal{T}$ is 2-DS (2-QS).

(ii) T_t is 2-DS (2-QS) for $t \in \mathbb{R}$ such that T_t is ergodic.

(iii) there exists $t_0 \in \mathbb{R}$ such that T_{t_0} is ergodic and T_{t_0} is 2-DS (2-QS).

Therefore, even though rather few explicit examples of 2-QS and 2-DS systems are known, these notions seem very natural. The obtained picture can be summarized as follows:



We present two proofs of Theorem 1.1. In the first approach we treat first the 2-QS property and then, building on the obtained results, we pass to the 2-DS property (see Section 3.3.1 and Section 3.3.2 up to page 27). The second approach yields an independent proof for the 2-DS property, see Section 3.3.2 from page 29 on. On the other hand, it does not provide much information about the 2-QS property. In both proofs, an essential role is played by lemmas, which may

be of an independent interest. They describe the relation between an extension of a flow and the corresponding extension of its time-one map. Roughly speaking, these extensions are of the same character (either both relatively distal or both relatively weakly mixing, see the definitions in Section 2.4).[7] In the first proof, our main tool is the so-called T-compactness property [47] (it describes isometric extensions). In the second proof we additionally use so-called separating sieves [63] (such sieves exist for distal extensions). For the definitions and for more details see Section 2.6. Even though it sounds now that the second proof, yielding a weaker result, is more complicated, this is not the case: by using separating sieves we can avoid some technical difficulties.

1.3.2 Mixing properties

In course of proving Theorem 1.1, as a by-product, we obtain a result which is related to mixing properties. Before we go into details, let us recall the following classical result:

Proposition 1.2. *Let $\mathcal{T}$ be a measurable flow. Then $\mathcal{T}$ is weakly mixing (mildly mixing, lightly mixing, partially mixing, mixing, mixing of order n) if and only if T_1 is weakly mixing (mildly mixing, lightly mixing, partially mixing, mixing, mixing of order n).*[8]

In the previous section, we already mentioned the property of relative weak mixing. It is therefore natural to expect that also a "relative version" of the above proposition is true. In course of proving Theorem 1.1, as a by-product, we will show that this is indeed the case as far as relative weak mixing is concerned. In fact, we will show more. In order to state the result, we need to recall first a classical result of Furstenberg and Zimmer:[9]

Theorem 1.3 (Furstenberg-Zimmer)**.** *Let T be an ergodic automorphism on $(X, \mathcal{B}, \mu)$ with a factor $\mathcal{A} \subset \mathcal{B}$. Then there exists a unique factor $\mathcal{C}$, such that $\mathcal{A} \subset \mathcal{C} \subset \mathcal{B}$ and such that the extension $T_1 \colon \mathcal{B} \to \mathcal{C}$ is relatively weakly mixing and $\mathcal{C} \to \mathcal{A}$ is relatively distal.*

The factorization from the above theorem is called *Furstenberg-Zimmer decomposition*. We show that for a flow and for its ergodic time-t automorphism this decomposition is the same with respect to any flow factor:[10]

Theorem 1.4. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}} \colon (X, \mathcal{B}, \mu) \to (X, \mathcal{B}, \mu)$ with T_1 ergodic and let $\mathcal{A} \subset \mathcal{B}$ be $\mathcal{T}$-invariant. Then the Furstenberg-Zimmer decomposition for $\mathcal{T}$ and for T_1 with respect to $\mathcal{A}$ is the same. In particular,*

- *$\mathcal{T}$ is relatively weakly mixing over $\mathcal{A}$ if and only if T_1 is relatively weakly mixing over $\mathcal{A}$,*
- *$\mathcal{T}$ is relatively distal over $\mathcal{A}$ if and only if T_1 is relatively distal over $\mathcal{A}$.*

[7] We are being a little bit imprecise, as we do not say here what is the invariant measure for the flow and for its time-one map.

[8] A similar result is true for rigidity and partial rigidity.

[9] The theorem is also valid for flows and other group actions.

[10] Again, an analogous result holds for automorphisms and their powers.

1.3.3 (Non-)uniqueness of the embedding

As already mentioned, Stepin and Eremenko [71] considered the problem of the uniqueness of the embedding. They showed that for T with simple spectrum, which is embeddable into a measurable flow, the embedding is either unique or there are infinitely many spectrally non-isomorphic flows with T as their time-one maps. In a typical situation the set of possible embeddings has cardinality continuum and any two different members are not spectrally isomorphic. Also the set of roots of a given order has typically cardinality of the continuum and any two different members of this set are not spectrally isomorphic.

In Section 4, we look at this problem from the point of view of self-joining properties. We show that for time-one map of a flow with minimal self-joinings the embedding is unique. A similar phenomenon can be observed for flows having the R-property, in particular for the horocycle flows. In both cases we also obtain the uniqueness of roots. For flows with the MSJ property we provide three independent proofs. One of them can be seen as a special case of a more general argument. Namely, we show that given a weakly mixing flow $\mathcal{T}$, a sufficient condition for the uniqueness of the embedding for T_1 is that any ergodic joining of $\mathcal{T}$ with a weakly mixing flow $\mathcal{S}$ remains weakly mixing. This is also the core of the argument we apply to the horocycle flows (using the theory of Ratner [64, 65, 66]).

Except for the Bernoulli shifts (both in finite and infinite entropy case) [55, 56, 57, 59], ergodic automorphisms being time-t maps of MSJ flows or flows with the R-property[11] seem to be the first concrete examples of automorphisms with a unique embedding into a measurable flow.

Finally, we pass to "negative" examples. They are given in terms of compact group extensions of 2-fold simple systems. To construct a weakly mixing 2-fold simple automorphism T which has two non-isomorphic square roots, we use the dihedral group of order 6 (one of these roots is 2-fold simple, the other one is only 2-QS). To give an example of a weakly mixing automorphism T which is 2-fold simple and embeddable into two non-isomorphic flows we use the group $SU(2)$ (again, one of the obtained flows is 2-fold simple, the other one is only 2-QS).[12] Summing up, the picture we have obtained is the following:

[11] Recall that both flows with the MSJ and R-property are of zero entropy.

[12] Contrary to what was claimed in [50], it is possible to have non-uniqueness of embedding and roots also in the quasi-discrete spectrum case, see also footnote 2. To see this, consider an irrational rotation on the circle $Tx = x + \alpha$. It has countably many non-isomorphic embeddings: $T_{n,t}x = x + t(\alpha + n)$. This yields immediately also countably many non-isomorphic roots for T. In a similar way, we can obtain non-isomorphic roots in the case of quasi-discrete, non-discrete spectrum. For $T(x, y) = (x+\alpha, x+y)$ we have $R \in C(T)$ for $R(x, y) = (x, y+1/2)$, whence $T^2 = (TR)^2$.



Another natural class of automorphisms, which is of a different nature,[13] with non-unique embedding into flows (and non-unique roots of any order) are Gaussian automorphisms.[14]

1.3.4 Counter-examples

In Section 5, we focus on various types of counter-examples related to Theorem 1.1. The examples mentioned at the end of the previous section show immediately that 2-fold simplicity of an (ergodic) time-one map of a flow does not imply that also the whole flow is 2-fold simple.[15] In other words, Theorem 1.1 is "the most" we can hope for, and the aforementioned result of del Junco and Rudolph [19] cannot be reversed. In fact, already Danilenko [11] constructed an automorphism T such that T^2 is 2-fold simple and T is only "2 : 1". The construction from [11] is highly non-trivial. He uses the so-called (C, F)-construction procedure, which is an algebraic counterpart of the well-known cutting and stacking technique.[16] A general idea is to carry out this type of construction using a larger (usually non-abelian) group with some special algebraic properties, which are reflected in the dynamics of a properly chosen sub-action. The example given by Danilenko actually has even stronger properties than the ones we were originally looking for: the constructed automorphism is 2-QS (it is "2:1"), it is disjoint from simple systems and its square is 2-fold simple. Notice that in the examples from the previous section each of the constructed actions (the automorphism T in the first example and the flow $\mathcal{T}$ in the second example) has a factor which is 2-fold simple. In particular, they are not disjoint from 2-fold simple systems. On the other hand, the construction of the automorphism from the previous section is much simpler than the example from [11].

Next we consider another example with the same basic properties: T^2 is 2-fold simple whereas T is only 2-QS. It is given in terms of a double group extension of an automorphism with the MSJ property (so it is again much simpler than the example from [11], but on the other hand not disjoint from

[13] It was shown in [45] that JP systems are disjoint from the Gaussian systems. Other related results can be found in [72] (disjointness of systems with the MSJ property and Gaussian systems), [17] (disjointness of simple systems from Gaussian systems) and [41] (disjointness of systems from some class generalizing the JP systems from Gausssian systems).

[14] We refer the reader e.g. to [10] or [53] for an introduction to Gaussian systems.

[15] Similarly, 2-fold simplicity of T^2 does not imply 2-fold simplicity of T.

[16] This method is a useful tool for producing examples and counter-examples of different types of behavior. For a detailed survey, see [12].

2-fold simple automorphisms). The second group extension in our construction is very explicit: we use the affine cocycle. This is however not necessarily a strong point: it is unclear how to adjust this example to obtain a 2-QS flow with a 2-fold simple time-one map.

In Section 5.2, using the construction from [11] as the starting point, we provide an example of a flow $\mathcal{T}$ with T_2 ergodic and 2-fold simple, such that $\mathcal{T}$ is only 2-QS (and not 2-fold simple). Moreover, the obtained flow has a factor with the same self-joining properties as the original flow and disjoint from 2-fold simple flows. To prove this, we show that given a 2-fold simple flow $\mathcal{T}\colon (X,\mathcal{B},\mu)\to(X,\mathcal{B},\mu)$ with T_t ergodic and a flow factor $\mathcal{A}\subset\mathcal{B}$, either both $\mathcal{T}|_{\mathcal{A}}$ and $T_t|_{\mathcal{A}}$ are 2-fold simple or none of them is. In particular, the horocycle flows and their factors cannot serve as an example here. The main difference between our (C, F)-construction and the constructions appearing previously in the literature is that we work in the setting of uncountable non-abelian groups which results in additional technical difficulties connected with equidistribution of sequences (see Section 6).

The full picture we have now looks like this:[17]



2 Preliminaries

In this paper we deal with measure-preserving automorphisms and flows, i.e. $\mathbb{Z}$- and $\mathbb{R}$-actions on standard Borel spaces, preserving a probability measure. In particular, we assume that all flows are measurable, i.e. the map $X\times\mathbb{R}\ni(x,t)\mapsto T_tx\in X$ is measurable. Sometimes we will deal with flows which are barely measurable, i.e. the invariant measure won't be fixed and we will have to do with a standard Borel space $(X,\mathcal{B})$. However, the existence of an invariant measure will be always guaranteed. The properties which are of our interest are invariant under measure-theoretical isomorphisms. Therefore we can assume without loss of generality that the flows we consider are continuous flows on compact metric spaces.[18] This will be our tacit assumption.

We will formulate most of the definitions in terms of $\mathbb{Z}$-actions. However, they can be transferred directly (or almost directly) to other group actions, in particular to $\mathbb{R}$-actions.

[17] See Question 1 on page 22.

[18] In the case of flows one uses the special flow representation [2, 3] and a result from [38] on the homology class of integrable functions.

2.1 Topology, invariant measures, ergodic decomposition

Let $(X, \mathcal{B})$ be a standard Borel space. Denote by $M_1(X)$ the set of probability measures on $(X, \mathcal{B})$. It is metrizable: for a dense set $\{f_i, i \geq 1\}$ of functions in $C(X)$, the metric given by

$$d(\mu_1, \mu_2) = \sum_{i \geq 1} \frac{1}{2^i \|f_i\|_{C(X)}} \left| \int_X f_i \, d\mu_1 - \int_X f_i \, d\mu_2 \right| \tag{1}$$

induces the weak topology which is compact.

Let $\mathcal{T} = (T_t)_{t \in \mathbb{R}}$ be measurable flow on a standard Borel space $(X, \mathcal{B})$. Let T also act measurably on $(X, \mathcal{B})$. Let

$$\begin{aligned}
&\mathcal{P}_{\mathbb{R}}(\mathcal{T}) = \{\mu \in M_1(X) \colon \mu \circ T_t = \mu \text{ for all } t \in \mathbb{R}\},\\
&\mathcal{P}^e_{\mathbb{R}}(\mathcal{T}) = \{\mu \in \mathcal{P}_{\mathbb{R}}(\mathcal{T}) \colon \mu \text{ is ergodic for } \mathcal{T}\},\\
&\mathcal{P}_{\mathbb{Z}}(T) = \{\mu \in M_1(X) \colon \mu \circ T = \mu\},\\
&\mathcal{P}^e_{\mathbb{Z}}(T) = \{\mu \in \mathcal{P}_{\mathbb{Z}}(T) \colon \mu \text{ is ergodic for } T\}.
\end{aligned}$$

Remark 2.1. Suppose that $\mathcal{P}_{\mathbb{R}}(\mathcal{T}) \neq \emptyset$. Then the map

$$\mathbb{R} \ni t \mapsto \mu \circ T_t \in M_1(X)$$

is continuous. Indeed, this follows easily by our assumption that $(X, \mathcal{B})$ is a compact metric space and $\mathcal{T}$ is continuous.

We will identify the following spaces: $\mathbb{R}/\mathbb{Z}$, $[0, 1)$, $\mathbb{T} = \{z \in \mathbb{C} \colon |z| = 1\}$ ($[0, 1)$ will be understood additively and $\mathbb{T}$ – multiplicatively) and we will equip them with the Lebesgue measure. Finally, given a measurable flow $\mathcal{T} = (T_t)_{t \in \mathbb{R}}$ and a measure $\nu \in \mathcal{P}^e_{\mathbb{Z}}(T_1)$ we set

$$\underline{\boldsymbol{\nu}} = \int_0^1 \nu \circ T_t \, dt. \tag{2}$$

Let us also recall the following folklore result (the proof can be found e.g. in [11]):

Lemma 2.1. *The measure $\underline{\boldsymbol{\nu}}$ is $\mathcal{T}$-invariant and ergodic. Moreover, the map*

$$\mathcal{P}^e_{\mathbb{Z}}(T_1) \ni \nu \mapsto \underline{\boldsymbol{\nu}} \in \mathcal{P}^e_{\mathbb{R}}(\mathcal{T}) \tag{3}$$

is onto and (2) is the ergodic decomposition of $\underline{\boldsymbol{\nu}}$ for T_1.

Since the measures $\nu \circ T_t$ in the decomposition (2) need not be pairwise orthogonal, the following lemma will be useful.

Lemma 2.2. *In the decomposition (2) either all measures $\nu \circ T_t$ are equal to ν or there exists $k \geq 1$ such that*

$$\underline{\boldsymbol{\nu}} = \int_{\mathbb{T}} \nu \circ T_t \, dt = k \int_0^{1/k} \nu \circ T_t \, dt \tag{4}$$

and for $0 \leq t < 1/k$ all measures $\eta \circ T_t$ are mutually singular.

Proof. Notice that any two measures in decomposition (2) are either equal or mutually singular. Let $A = \{t \in \mathbb{T}\colon \eta = \eta \circ T_t\}$. It is clearly a subgroup of $\mathbb{T}$ which is closed by Remark 2.1. Therefore either $A = \mathbb{T}$ or A is finite. In the latter case we obtain formula (4) with $k = \#A$. □

Remark 2.2. If $\mathcal{T}$ is a suspension flow with the first return time to the base equal to $a > 0$ and a weakly mixing first return time map, then the discrete part of the spectrum is concentrated on $\frac{1}{a}\mathbb{Z}$.

Remark 2.3. Recall that a flow is a suspension flow (with a constant first return time to the base) if and only it has a rational point in its discrete spectrum [79]. Therefore, whenever (4) holds, the flow $\mathcal{T}$ can be represented as a suspension flow over its $1/k$-time map with common return time to the base equal to $1/k$. In other words, the whole space can be identified with $\big(X \times [0,1/k], \nu \otimes \lambda_{[0,1/k]}\big)$, with the action of $(T_t)_{t\in\mathbb{R}}$ given by $T_t(x,s) = (x, s+t)$ and the identification of the points $(x, 1/k)$ and $(T_{1/k}x, 0)$. Moreover, by Remark 2.2, the discrete part of its spectrum is concentrated on $k\mathbb{Z}$.

2.2 Factors and extensions I

Let $T\colon (X,\mathcal{B},\mu) \to (X,\mathcal{B},\mu)$ and $S\colon (Y,\mathcal{C},\nu) \to (Y,\mathcal{C},\nu)$ be automorphisms of standard probability Borel spaces. We say that S is a *factor* of T (T is an *extension* of S) if there exists $\pi\colon X \to Y$ such that $\pi \circ T = S \circ \pi$ and $\nu = \pi_*(\mu)$, i.e. for any $A \in \mathcal{C}$ we have $\mu(\pi^{-1}A) = \nu(A)$. We write $T \to S$. Notice that when S is a factor of T then $\pi^{-1}(\mathcal{C}) \subset \mathcal{B}$ is T-invariant. On the other hand, every T-invariant sub-σ-algebra $\mathcal{A} \subset \mathcal{B}$ corresponds to some factor of T. If no confusion arises we will often write $\mathcal{B} \to \mathcal{A}$.

2.3 Joinings I

2.3.1 Basic definitions and properties

Recall that $T\colon (X,\mathcal{B},\mu) \to (X,\mathcal{B},\mu)$ and $S\colon (Y,\mathcal{C},\nu) \to (Y,\mathcal{C},\nu)$ are automorphisms of standard probability Borel spaces. By $\mathcal{J}(T,S)$ we denote the set of all *joinings* between T and S, i.e. the set of all $T \times S$-invariant probability measures on $(X \times Y, \mathcal{B} \otimes \mathcal{C})$, whose projections on X and Y are equal to μ and ν respectively. For $\mathcal{J}(T,T)$ we write $\mathcal{J}(T)$ and we speak about *self-joinings*. If we assume additionally that both T and S are ergodic, then $\mathcal{J}(T,S)$ is a simplex whose extreme points are joinings which are ergodic with respect to $T \times S$. We denote the set of such joinings by $\mathcal{J}^e(T,S)$ (or $\mathcal{J}^e(T)$ if $S = T$). If $\mathcal{J}(T,S) = \{\mu \otimes \nu\}$ we say that T and S are disjoint [26] and we write $T \perp S$.

In a similar way one defines joinings of higher order, i.e. joinings of more than two automorphisms. We will denote them by $\mathcal{J}(T_1,\dots,T_k)$, $k \geq 2$ and $\mathcal{J}(T_1,T_2,\dots)$ in case of a finite and an infinite number of automorphisms respectively. Again, we may be interested in the situation where all T_i are isomorphic and speak about self-joinings. To distinguish self-joinings of different orders we write e.g. $\mathcal{J}^e_\infty(T)$ for infinite ergodic self-joinings of T or $\mathcal{J}_3(T)$ for three-fold self-joinings of T. By $\mathcal{J}_2(T;\mathcal{A})$, where $\mathcal{A}$ is a factor of T, we denote all two-fold self-joinings of T which project onto $\mathcal{A}\otimes\mathcal{A}$ as the diagonal measure. Sometimes, to simplify the notation, we will denote a joining between T and S by $T \vee S$.

Recall that 2-fold joinings are in one-to-one correspondence with Markov operators

$$\Phi\colon L^2(X,\mathcal{B},\mu)\to L^2(Y,\mathcal{C},\nu)$$

such that $\Phi\circ T = S\circ\Phi$. This identification allows us to view $\mathcal{J}_2(T)$ as a metrisable compact semitopological semigroup endowed with the weak operator topology.

The metric inducing the weak topology on $\mathcal{J}_2(T)$ can be defined in the following way. Fix a dense subset $\{f_i'\colon i\geq 1\}\subset L^2(X,\mathcal{B},\mu)$ and put $f_i := f_i'/\|f_i'\|_{L^2(X,\mu)}$ for $i\geq 1$. Define a metric $\overline{d}_{\{f_n\}_{n\in\mathbb{N}}}$ by the formula

$$\begin{aligned}&\overline{d}_{\{f_n\}_{n\in\mathbb{N}}}(\xi,\nu)\\ &\quad=\sum_{i,j=1}^{\infty}\frac{1}{2^{i+j}}\left|\int_{X\times X} f_i(x)\overline{f_j(y)}\,d\xi(x,y)-\int_{X\times X} f_i(x)\overline{f_j(y)}\,d\nu(x,y)\right|.\end{aligned}\tag{5}$$

Notice that for $\xi,\nu\in\mathcal{J}_2(T)$ we always have

$$\overline{d}_{\{f_n\}_{n\in\mathbb{N}}}(\xi\circ(T\times T),\nu\circ(T\times T))=\overline{d}_{(f_n)_{n\in\mathbb{N}}}(\xi,\nu).\tag{6}$$

Remark 2.4. This metric is equivalent to the metric d introduced in (1). Instead of representing X as a compact metric space and taking continuous functions on $X\times X$, we take tensors of the form $f_i\otimes f_j$, with $f_i,f_j\in L^2(X,\mu)$. This is more convenient as joinings are measures defined on the product σ-algebra and the marginals are fixed.

Remark 2.5. We will often use Lemma 2.1 and Lemma 2.2 for joinings. Notice that given a flow $\mathcal{T}$ with T_1 ergodic, the map $\eta\mapsto\underline{\boldsymbol{\eta}}$ has values in $\mathcal{J}_2^e(\mathcal{T})$ whenever $\nu\in\mathcal{J}_2^e(T_1)$. Moreover, it is surjective.

Basic types of self-joinings

- *The product* measure $\mu\otimes\mu$ is always a self-joining.
- For $S_2,\ldots,S_N\in C(T)$[19] the measure $\nu=\mu_{S_2,\ldots,S_N}$ defined by

  $$\mu_{S_2,\ldots,S_N}(A_1\times\cdots\times A_N)=\mu(A_1\cap S_2^{-1}A_2\cap\cdots\cap S_N^{-1}(A_N))$$

  (with an obvious modification for $N=\infty$) is an N-self-joining of T called a *graph self-joining* (or *off-diagonal self-joining*). If $S_2=\cdots=S_N=Id$, we speak about the *diagonal self-joining* and we denote it by Δ_N. For $N=2$ we often write Δ instead of Δ_2. Sometimes, when a confusion might arise, we denote the marginals of Δ in parenthesis: $\Delta(\mu)$.
- Each pair of factors $\mathcal{A}_1,\mathcal{A}_2\subset\mathcal{B}$, together with $\lambda\in\mathcal{J}(T|_{\mathcal{A}_1},T|_{\mathcal{A}_2})$ yield a self-joining of T defined by

  $$\widehat{\lambda}(A_1\times A_2)=\int_{X/\mathcal{A}_1\times X/\mathcal{A}_2}E(A_1|\mathcal{A}_1)(\overline{x})E(A_2|\mathcal{A}_2)(\overline{y})\,d\lambda(\overline{x},\overline{y}).$$

  It is called a *relatively independent extension of* λ. In particular, when $\mathcal{A}_1=\mathcal{A}_2=\mathcal{A}$ and $\lambda=\Delta$ we say that $\widehat{\Delta}$ is a relatively independent extension over $\mathcal{A}$. We denote it by $\mu\otimes_{\mathcal{A}}\mu$.

[19] $C(T)$ stands for the centralizer of T, i.e. the group of all measure-preserving automorphisms which commute with T.

- Let $\lambda \in \mathcal{J}(T_1, \dots, T_n)$. Then $\lambda|_{X_i,X_j} = \lambda|_{X_i \times X_j}$ defined by

  $$\nu|_{X_i \times X_j}(C \times D) = \nu(\underbrace{X \times \cdots \times X}_{i-1} \times C \times \underbrace{X \times \cdots \times X}_{j-i-1} \times D \times \underbrace{X \times \cdots \times X}_{n-j})$$

  is an element of $\mathcal{J}(T_i, T_j)$. Moreover, if $\lambda \in \mathcal{J}^e(T_1, \dots, T_n)$ then $\lambda|_{X_i,X_j} \in \mathcal{J}^e(T_i, T_j)$.

- Let $\lambda \in \mathcal{J}(T_1, T_2)$ and let $\mathcal{A}_1, \mathcal{A}_2$ be factors of T_1, T_2, respectively. Then $\lambda|_{\mathcal{A}\otimes\mathcal{C}} \in \mathcal{J}((T_1)|_{\mathcal{A}}, (T_2)|_{\mathcal{C}})$.

The above mentioned examples of self-joinings can be easily modified to obtain joinings of two or more different automorphisms.

2.4 Factors and extensions II

Any ergodic extension $\widetilde{T} \to T$ is isomorphic to a *skew product* over T, i.e. to an automorphism $\overline{T}\colon (X \times Y, \mathcal{B} \otimes \mathcal{C}, \mu \otimes \nu) \to (X \times Y, \mathcal{B} \otimes \mathcal{C}, \mu \otimes \nu)$ which is of the form

$$\overline{T}(x, y) = (Tx, S_x y), \tag{7}$$

where $(S_x)_{x\in X}$ is a measurable family of automorphisms of some standard probability space $(Y, \mathcal{C}, \nu)$ [1]. A particular case of skew products are *Rokhlin extensions*, i.e. automorphisms of the form

$$T_{S,\varphi}(x, y) = (Tx, S_{\varphi(x)} y),$$

where $(S_g)_{g\in G}$ is a measurable representation of a locally compact abelian group G in the group of automorphisms of $(Y, \mathcal{C}, \nu)$ and $\varphi\colon X \to G$ is a measurable function.

2.4.1 Relatively weakly mixing extensions

Following [27], we say that $T\colon (X, \mathcal{B}, \mu) \to (X, \mathcal{B}, \mu)$ is *relatively weakly mixing* with respect to factor $\mathcal{A} \subset \mathcal{B}$ if the relatively independent extension of the diagonal self-joining of $T|_{\mathcal{A}}$ is ergodic. In terms of the skew product representation (7), the property of relative weak mixing is equivalent to ergodicity of automorphism $\overline{\overline{T}}$ acting on $(X \times Y \times Y, \mathcal{B} \otimes \mathcal{C} \otimes \mathcal{C}, \mu \otimes \nu \otimes \nu)$, given by the formula $\overline{\overline{T}}(x, y_1, y_2) = (Tx, S_x y_1, S_x y_2)$.

2.4.2 Relatively isometric and relatively distal extensions

Relative distality is a concept which is of "opposite nature" to relative weak mixing (cf. Theorem 1.3 on page 6 on the Furstenberg-Zimmer decomposition).

Compact group extensions Let us first recall the definition of a compact group extension. Let G be a compact metrizable group with the normalized Haar measure λ_G and let $\varphi\colon X \to G$ be measurable. Then $T_\varphi\colon (x, g) \mapsto (Tx, \varphi(x)g)$ is an automorphism of $(X \times G, \mathcal{B} \otimes \mathcal{B}(G), \mu \otimes \lambda_G)$. We call such an automorphism a (compact) *group extension* of T. Compact group extensions can be characterized in terms of self-joinings:

Proposition 2.3 (see e.g.[19]). *If $T\colon (X,\mathcal{B},\mu)\to(X,\mathcal{B},\mu)$ is an ergodic compact group extension of its factor $\mathcal{A}$ then $J_2^e(T;\mathcal{A})$ consists of graphs joinings.*

Proposition 2.4 ($\mathbb{Z}$-actions: [72, 75], actions of other groups: [19]). *If $T\colon (X,\mathcal{B},\mu)\to(X,\mathcal{B},\mu)$ is ergodic, $\mathcal{A}$ is a factor of T and $J_2^e(T;\mathcal{A})$ consists of graphs joinings then $\mathcal{B}\to\mathcal{A}$ is a compact group extension.*

We will also use the following fact:

Lemma 2.5. *[43] Let $T\colon (X,\mathcal{B},\mu)\to(X,\mathcal{B},\mu)$ be a (not necessarily ergodic) compact group extension of an ergodic factor $\mathcal{A}$. Then every ergodic component of T is also isomorphic to some compact group extension of $\mathcal{A}$.*

Isometric extensions Let G be a compact group with a closed subgroup H and let $\varphi\colon X\to G$ be measurable. The action of T_φ restricted to $X\times G/H$ is called an *isometric extension* of T.

In [78] Zimmer introduced the notion of relatively discrete spectrum. It is based on the classical notion of discrete spectrum of a unitary representation. As we will not need the precise definition, let us remark here only that

an ergodic extension has relatively discrete spectrum
if and only if it is an isometric extension. (8)

Distal extensions We say that the extension $\widetilde{T}\colon (\widetilde{X},\widetilde{\mathcal{B}},\widetilde{\mu})\to(\widetilde{X},\widetilde{\mathcal{B}},\widetilde{\mu})$ of $T\colon (X,\mathcal{B},\mu)\to(X,\mathcal{B},\mu)$ is *relatively distal* [77] if there exists a transfinite sequence $(\mathcal{B}_\alpha)_{\alpha\le\beta}$ of factors of $\widetilde{T}$ such that $\mathcal{B}_0=\mathcal{B}$, $\mathcal{B}=\widetilde{\mathcal{B}}$, the extensions $\mathcal{B}_{\alpha+1}\to\mathcal{B}_\alpha$ are isometric and for α being limit ordinals, $\mathcal{B}_\alpha$ is the inverse limit of the preceding factors.

2.5 Joinings II

2.5.1 Self-joining properties

Let $T\colon (X,\mathcal{B},\mu)\to(X,\mathcal{B},\mu)$ be an ergodic automorphism. It is said that T

- has the *pairwise independence property* (PID) if every (finite or infinite) pairwise independent self-joining of T is equal to the product measure.[20]

Moreover, it is said that T

- has *minimal self-joinings* if the only ergodic self-joinings of T (other than the product measure) are off-diagonal and $C(T)=\{T^k\colon k\in\mathbb{Z}\}$;
- is *2-fold simple* if the only ergodic self-joinings of T (other than the product measure) are off-diagonal;
- is *simple* if it is 2-fold simple and PID;
- is *2-fold quasi-simple* (2-QS) if the only ergodic self-joinings of T (other than the product measure) are isometric extensions of each of the coordinate factors;

[20] The PID property was introduced in [19]. For the references to other notions recalled below, see Section 1.2.

- is *quasi-simple* if it is 2-QS and PID;
- is *"n:1"* for some $n \in \mathbb{N}$ if the only ergodic self-joinings of T (other than the product measure) are finite "$n:1$"-extensions of each of the coordinate factors (this is a special case of 2-QS);
- is *2-fold distally-simple* (2-DS) if the only ergodic self-joinings of T (other than the product measure) are distal extensions of each of the coordinate factors.

The next property is the *joining primeness property* (JP). The original definition refers to joinings with other automorphisms, however, we will show later that there is an equivalent intrinsic definition, i.e. one given in terms of self-joinings. We say that T has joining primeness property if for every pair of weakly mixing systems S_1, S_2 and every $\lambda \in \mathcal{J}^e(R, S_1 \times S_2)$ we have

$$\lambda = \lambda|_{Z,Y_1} \otimes \nu_2 \text{ or } \lambda = \lambda|_{Z,Y_2} \otimes \nu_1.$$

Notice that assuming S_1 and S_2 isomorphic would give an equivalent definition.

The last property which we recall here is the so-called *Ratner property* (R-property). It is a property which can be enjoyed by flows and it has a different flavor than all the properties mentioned above. Since the definition is very technical and we will not use it explicitly, let us list here some examples of flows which have this property and refer the reader e.g. to [25] for more details:

- horocycle flows [66],
- special flows [21] over the rotations by an irrational α with bounded partial quotients under f which is piecewise absolutely continuous with a non-zero sum of jumps [24],
- special flows over two-dimensional rotations by (α, β) on $\mathbb{T}^2$ under piecewise C^2 roof functions f satisfying von Neumann condition

$$\int_{\mathbb{T}^2} f_x(x,y)\ dx\ dy \neq 0 \neq \int_{\mathbb{T}^2} f_y(x,y)\ dx\ dy \text{ [25]},$$

- special flows over the rotations by an irrational α under the roof functions of bounded variation without continuous, singular part in the Lebesgue decomposition and a non-zero sum of jumps [35].

Let us remark that the definition of R-property itself does not refer to joinings at all, however it imposes restrictions on the joinings of the flow under consideration. The following proposition is a little bit imprecise (as we don't say what we mean by the R-property here), but it will be sufficient for our needs in this form. Moreover, the aforementioned examples do satisfy all the assumptions.

Proposition 2.6 ([25]). *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be a weakly mixing flow on $(X, \mathcal{B}, \mu)$ which has the R-property. Let $\mathcal{S} = (S_t)_{t\in\mathbb{R}}$ be an ergodic flow on $(Y, \mathcal{C}, \nu)$ and let $\lambda \in \mathcal{J}^e(\mathcal{T}, \mathcal{S})$. Then either $\lambda = \mu \otimes \nu$ or λ is a finite extension of ν.*

[21] As we will not need the notion of a special flow for stating or proving the results, we do not recall it here. Instead, we refer the reader e.g. to [10].

2.6 Distal and isometric extensions – tools

2.6.1 Separating sieves

Let $T\colon (X,\mathcal{B},\mu)\to(X,\mathcal{B},\mu)$ and $S\colon (Y,\mathcal{C},\nu)\to(Y,\mathcal{C},\nu)$ be automorphisms of standard probability Borel spaces. Suppose that T is an ergodic extension of S. Let $\pi\colon X\to Y$ be the factoring map.

Definition 2.1 ([63])**.** Given $A\in\mathcal{B}$, let

$$A\times_Y A=\{(x,x')\in A\times A\colon \pi(x)=\pi(x')\}.$$

A sequence $A_1\supset A_2\supset\dots$ of sets in $\mathcal{B}$ with $\mu(A_n)>0$ and $\mu(A_n)\to 0$, is called a *separating sieve* over Y if there exists a subset $X_0\subset X$ with $\mu(X_0)=1$ such that for every $x,x'\in X_0$ with $\pi(x)=\pi(x')$, the condition "for every $n\in\mathbb{N}$ there exists $m\in\mathbb{Z}$ with $T^mx,T^mx'\in A_n$" implies $x=x'$, i.e.

$$\bigcap_{n=1}^{\infty}\left(\bigcup_{m\in\mathbb{Z}}(A_n\times_Y A_n)\right)\cap(X_0\times_Y X_0)\subset\{(x,x)\colon x\in X_0\}.$$

We also say that the extension π has a *separating sieve*.

Remark 2.6. Zimmer [78] showed that the existence of a separating sieve is equivalent to distality of the considered extension.

Remark 2.7. Notice that whenever $(A_n)_{n\in\mathbb{N}}$ forms a separating sieve for the extension π, then any subsequence $(A_{n_k})_{k\geq 1}$ is a separating sieve for π. Also, any decreasing sequence $(B_n)_{n\in\mathbb{N}}$ of sets of positive measure with $B_n\subset A_n$ forms a separating sieve for π.

Lemma 2.7. [22] *Suppose that T is an ergodic extension of S with the factoring map π. The following conditions are equivalent:*

(i) There exists a separating sieve for π.

(ii) There exists $X_0\subset X$ with $\mu(X_0)=1$ and a sequence of sets $(B_n)_{n\in\mathbb{N}}$ with $\mu(B_n)>0$, $\mu(B_n)\to 0$ and such that for $x,x'\in X$ with $\pi(x)=\pi(x')$ whenever for every $n\in\mathbb{N}$ there exists $m_n\in\mathbb{Z}$ such that $T^{m_n}x,T^{m_n}x'\in B_n$ then $x=x'$.

Moreover, if additionally μ is non-atomic, the above conditions are equivalent to

(ii') There exists $X_0\subset X$ with $\mu(X_0)=1$ and a sequence of sets $(B_n)_{n\in\mathbb{N}}$, $\mu(B_n)>0$ and such that for $x,x'\in X$ with $\pi(x)=\pi(x')$ whenever for every $n\in\mathbb{N}$ there exists $m_n\in\mathbb{Z}$ such that $T^{m_n}x,T^{m_n}x'\in B_n$ then $x=x'$.

Proof. We will show first that (ii) implies (i). Let us recursively define a sequence of sets $(A_n)_{n\in\mathbb{N}}$:

$$A_1=B_1,\ A_n=T^{k_n}B_n\cap A_{n-1},$$

[22] An analogous result holds for other group actions – the proof can be repeated word by word.

where $k_n \in \mathbb{Z}$ is such that $\mu(A_n) = \mu(T^{k_n} B_n \cap A_{n-1}) > 0$. Such k_n exists by ergodicity of T. We claim that $(A_n)_{n\in\mathbb{N}}$ is a separating sieve for π. Clearly $\mu(A_n) \to 0$. Let $x, x' \in X_0$ be such that for each $n \in \mathbb{N}$ there exists m_n such that $T^{m_n} x, T^{m_n} x' \in A_n$. By (ii) this implies that $T^{-k_n+m_n} x, T^{-k_n+m_n} x' \in B_n$ whence $x = x'$ and we have proved that (i) follows from (ii).

To see that (ii') implies (i), it suffices to modify the proof of the previous part by adding one more step. Namely, we make the sets A_n smaller, if necessary, in such a way that $\mu(A_n) \to 0$.

Clearly (i) implies (ii) and (ii'). □

2.6.2 T-compactness

Let $\mathcal{A}$ be a factor of T. Following [47], we say that the extension $\mathcal{B} \to \mathcal{A}$ is *T-compact* if for any $\varepsilon > 0$ there exists $N \geq 2$ such that for each $\nu \in J^e_N(T; \mathcal{A})$ we can find $1 \leq i < j \leq N$ such that

$$\overline{d}(\nu|_{X_i \times X_j}, \Delta_N|_{X_i \times X_j}) < \varepsilon.^{23}$$

Proposition 2.8 ([47]). *Assume that $T\colon (X, \mathcal{B}, \mu) \to (X, \mathcal{B}, \mu)$ is an ergodic automorphism and let $\mathcal{A} \subset \mathcal{B}$ be its factor. Then the following conditions are equivalent:*

(i) $\mathcal{B} \to \mathcal{A}$ has relatively discrete spectrum,

(ii) $\mathcal{B} \to \mathcal{A}$ is isometric,

(iii) $\mathcal{B} \to \mathcal{A}$ is T-compact.

3 Self-joining properties of $\mathcal{T}$ and T_1 – part I

3.1 Simplicity

Our starting point is a result of del Junco and Rudolph from [19]. We state it in a slightly different form than the original one and also include its proof. The language we use in the proof is compatible with the language used later on.

Proposition 3.1. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be a weakly mixing flow. Then $\mathcal{T}$ is 2-fold simple if and only if T_1 is 2-fold simple and $C(\mathcal{T}) = C(T_1)$.*[24]

Remarks

1. Notice that in particular, $C(\mathcal{T}) = C(T_1)$ whenever $C(T_1)$ is abelian.
2. An analogous result to Proposition 3.1 holds for automorphisms and their powers.
3. A flow $\mathcal{T}$ which is not weakly mixing and which has an ergodic time T_t which is 2-fold simple, is also 2-fold simple. Indeed, as $\mathcal{T}$ is not weakly mixing, T_t is also not weakly mixing, whence it has purely discrete spectrum [19]. Therefore the whole flow $\mathcal{T}$ has purely discrete spectrum and it follows that it is 2-fold simple.

[23] See the definitions in Section 2.3.

[24] Del Junco and Rudolph showed [19] that if $\mathcal{T}$ is 2-fold simple then T_1 is also 2-fold simple and $C(\mathcal{T}) = C(T_1)$.

Proof of Proposition 3.1. Suppose that $\mathcal{T}$ is 2-fold simple. We want to show that $\mathcal{J}_2^e(\mathcal{T}) = \mathcal{J}_2^e(T_1)$. It will follow immediately that T_1 is also 2-fold simple and $C(\mathcal{T}) = C(T_1)$, as every ergodic 2-fold self-joining for $\mathcal{T}$, other than the product measure, corresponds to an element $S \in C(\mathcal{T})$. By the assumption, we have $\mathcal{J}_2^e(\mathcal{T}) \subset \mathcal{J}_2^e(T_1)$. Suppose that we can find $\eta \in \mathcal{J}_2^e(T_1) \setminus \mathcal{J}_2^e(\mathcal{T})$. Define

$$\underline{\boldsymbol{\eta}} = \int_0^1 \eta \circ (T_t \times T_t)\, dt. \tag{9}$$

By Remark 2.5 on page 12, we have $\underline{\boldsymbol{\eta}} \in \mathcal{J}_2^e(\mathcal{T}) \subset \mathcal{J}_2^e(T_1)$. It follows by Lemma 2.2 and by the uniqueness of ergodic decomposition that the decomposition (9) is trivial, i.e. we have $\underline{\boldsymbol{\eta}} = \eta$, which yields a contradiction.

Suppose now that T_1 is 2-fold simple and $C(\mathcal{T}) = C(T_1)$. Therefore $\mathcal{J}_2^e(T_1) \subset \mathcal{J}_2^e(\mathcal{T})$, whence $\mathcal{J}_2(T_1) \subset \mathcal{J}_2(\mathcal{T})$. The opposite inclusion $\mathcal{J}_2(\mathcal{T}) \subset \mathcal{J}(T_1)$ is obvious and the claim follows.

□

Remark 3.1. Let T and S be 2-fold simple automorphisms such that $T^k = S^l$ for some $k, l \in \mathbb{Z} \setminus \{0\}$. It follows from Proposition 3.1 that $T^k = S^l$ is 2-fold simple and $C(T) = C(T^k) = C(S^l) = C(S)$. In particular, $\mathcal{J}_2^e(T) = \mathcal{J}_2^e(S)$.

For a measure-preserving flow $\mathcal{T}$ let

$$I_s(\mathcal{T}) = \{t \in \mathbb{R} \colon T_t \text{ is 2-fold simple}\} \cup \{0\}.$$

Lemma 3.2. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be a weakly mixing flow. Suppose that T_α and T_β are 2-fold simple for some $\alpha, \beta \neq 0$ such that $\alpha/\beta \notin \mathbb{Q}$. Then $\mathcal{T}$ is 2-fold simple. In particular, $I_s(\mathcal{T}) = \mathbb{R}$.*

Proof. We may assume without loss of generality that $\beta = 1$ and $\alpha \notin \mathbb{Q}$. There are two possibilities:

(i) $C(T_1) = C(T_\alpha)$,

(ii) $C(T_1) \neq C(T_\alpha)$.

We first cover case (i). Fix $t_0 \in \mathbb{R}$. For $n \in \mathbb{N}$ choose $t_n \in \mathbb{Z} + \alpha\mathbb{Z}$ such that $t_n \to t_0$ (such a choice can always be made as $\alpha \notin \mathbb{Q}$). Take $S \in C(T_1) = C(T_\alpha)$. We have

$$ST_{t_n} \to ST_{t_0} \text{ and } T_{t_n}S \to T_{t_0}S.$$

Since $ST_{t_n} = T_{t_n}S$, it follows that $ST_{t_0} = T_{t_0}S$, i.e. $C(T_1) = C(\mathcal{T})$. By Proposition 3.1, $\mathcal{T}$ is 2-fold simple.

We pass now to case (ii). We may assume without loss of generality that there exists $S \in C(T_1) \setminus C(T_\alpha)$. Consider

$$\underline{\boldsymbol{\mu_S}} := \int_0^1 \mu_S \circ (T_t \times T_t)\, dt. \tag{10}$$

By Remark 2.5, $\underline{\boldsymbol{\mu_S}} \in \mathcal{J}_2^e(\mathcal{T})$. Clearly $\underline{\boldsymbol{\mu_S}} \notin J_2^e(T_1)$ (otherwise the measures in the decomposition (10) would be all equal, which would imply that $S \in C(\mathcal{T}) \subset C(T_\alpha)$). By Lemma 2.2 the ergodic decomposition of $\underline{\boldsymbol{\mu_S}}$ for $T_1 \times T_1$ is of the form

$$\underline{\boldsymbol{\mu_S}} = k \int_0^{1/k} \mu_S \circ (T_t \times T_t)\, dt \tag{11}$$

for some $k \geq 1$. By Remark 2.3

$$\text{the set of eigenvalues of } (\mathcal{T} \times \mathcal{T}, \underline{\boldsymbol{\mu}_{\boldsymbol{S}}}) \text{ is equal to } k\mathbb{Z}. \tag{12}$$

Suppose that $\underline{\boldsymbol{\mu}_{\boldsymbol{S}}}$ is not ergodic for $T_\alpha \times T_\alpha$. Then by Lemma 2.1

$$\underline{\boldsymbol{\mu}_{\boldsymbol{S}}} = \frac{1}{\alpha} \int_0^\alpha \mu_W \circ (T_t \times T_t)\, dt$$

for some $W \in C(T_\alpha)$. By Lemma 2.2 the ergodic decomposition of $\underline{\boldsymbol{\mu}_{\boldsymbol{S}}}$ for $T_\alpha \times T_\alpha$ is therefore of the form

$$\underline{\boldsymbol{\mu}_{\boldsymbol{S}}} = \frac{l}{\alpha} \int_0^{\alpha/l} \mu_W \circ (T_t \times T_t)\, dt \tag{13}$$

for some $l \geq 1$. By Remark 2.3 the set of eigenvalues $(\mathcal{T} \times \mathcal{T}, \underline{\boldsymbol{\mu}_{\boldsymbol{S}}})$ is equal to $(l/\alpha)\mathbb{Z}$. This yields a contradiction with (12).

Suppose now that $\underline{\boldsymbol{\mu}_{\boldsymbol{S}}}$ is ergodic for $T_\alpha \times T_\alpha$. Then $\underline{\boldsymbol{\mu}_{\boldsymbol{S}}} = \mu_W$ for some $W \in C(T_\alpha)$. It follows that $(T_\alpha \times T_\alpha, \underline{\boldsymbol{\mu}_{\boldsymbol{S}}})$ is weakly mixing, whence also $(\mathcal{T} \times \mathcal{T}, \underline{\boldsymbol{\mu}_{\boldsymbol{S}}})$ is weakly mixing. This yields a contradiction with (12) and ends the proof. □

Proposition 3.3. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be a weakly mixing flow. Then $I_s(\mathcal{T}) = \mathbb{R}$ or $I_s(\mathcal{T}) = \beta\mathbb{Z}$ for some $\beta \in \mathbb{R}$. Moreover, if $I_s(\mathcal{T}) = \mathbb{R}$ then $\mathcal{T}$ is 2-fold simple.*

Proof. Suppose that $I_s(\mathcal{T}) \neq \beta\mathbb{Z}$ for all $\beta \in \mathbb{R}$. Consider the following two possibilities:

(i) there exist $t_1, t_2 \in I_s(\mathcal{T}) \setminus \{0\}$ which are rationally independent,

(ii) all numbers in $I_s(\mathcal{T})$ are rationally dependent.

In case (i) it follows from Lemma 3.2 that $\mathcal{T}$ is 2-fold simple and $I_s(\mathbb{R}) = \mathbb{R}\setminus\{0\}$. In case (ii) there exist $\alpha > 0$ such that $I_s(\mathcal{T}) \subset \alpha\mathbb{Q}$. Take $t_1, t_2 \in I_s(\mathcal{T})$. We claim that the whole additive subgroup generated by t_1 and t_2 is in $I_s(\mathcal{T})$. Indeed, we have $t_1 = p_1/q$, $t_2 = p_2/q$ for some $p_1, p_2, q \in \mathbb{Z}$. Let $d = \gcd(p_1, p_2)$. Then for some $k, l \in \mathbb{Z}$ we have $kp_1 + lp_2 = d$. Hence $d/q = kp_1/q + lp_2/q$. It follows by Remark 3.1 that $C(T_{p_1/q}) = C(T_{p_2/q})$. Therefore $C(T_{p_1/q}) \subset C(T_{d/q})$. On the other hand, $C(T_{d/q}) \subset C(T_{p_1/q})$ since $T_{d/q}$ is a root of $T_{p_1/q}$. Hence $C(T_{d/q}) = C(T_{p_1/q})$ and it follows that $T_{d/q}$ is 2-fold simple. Therefore, for all $k \in \mathbb{Z}$ also $T_{kd/q}$ is 2-fold simple. It follows that $I_s(\mathcal{T}) = I_s(\mathcal{T}) \cap \alpha\mathbb{Q}$ is an additive subgroup of $\alpha\mathbb{Q}$. Notice that $I_s(\mathcal{T})$ cannot be dense in $\mathbb{R}$. Indeed, if it was dense, an argument similar to the one we used in the proof of (i) of Lemma 3.2 would imply $I_s(\mathcal{T}) = \mathbb{R}$. The claim follows. □

3.2 Joining primeness property

A natural approach is to try to extend Proposition 3.1 to other classes of dynamical systems listed in Section 2.5.1. It turns out that a converse of Proposition 3.1 is true if we replace "2-fold simple" with "JP":

Proposition 3.4. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be a measure-preserving flow with T_1 ergodic. If T_1 has the JP property then $\mathcal{T}$ has the JP property.*

Proof. Let $\underline{\boldsymbol{\eta}} \in J^e(\mathcal{T}, \mathcal{R} \times \mathcal{S}) \subset J(T_1, R_1 \times S_1)$ for some weakly mixing flows $\mathcal{R}$ and $\mathcal{S}$. If $\underline{\boldsymbol{\eta}} \in J^e(T_1, R_1 \times S_1)$ then $\underline{\boldsymbol{\eta}} = \underline{\boldsymbol{\eta}}|_{X\times Y} \otimes \nu_2$ or $\underline{\boldsymbol{\eta}} = \underline{\boldsymbol{\eta}}|_{X\times Z} \otimes \nu_1$. If $\underline{\boldsymbol{\eta}} \notin J^e(T_1, R_1 \times S_1)$ then by Lemma 2.1, Lemma 2.2 and Remark 2.5 on page 12, we may assume without loss of generality that for some $\eta \in J^e(T_1, R_1 \times S_1)$ we have

$$\underline{\boldsymbol{\eta}} = \int_0^1 \eta \circ (T_t \times R_t \times S_t)\, dt.$$

Notice that η cannot be the product measure. Indeed, it would be invariant under the product flow $\mathcal{R} \times \mathcal{S} = (R_t \times S_t)_{t\in\mathbb{R}}$ and equal to $\underline{\boldsymbol{\eta}}$, which is impossible since $\underline{\boldsymbol{\eta}} \notin \mathcal{J}^e(T_1, R_1 \times S_1)$. Therefore, up to a permutation of the names of the coordinates, we have

$$\eta = \eta|_{X\times Y} \otimes \nu_2.$$

Hence

$$\underline{\boldsymbol{\eta}} = \int_0^1 \eta|_{X\times Y} \circ (T_t \times R_t) \otimes \nu_2 \, dt = \left(\int_0^1 \eta|_{X\times Y} \circ (T_t \times R_t)\, dt \right) \otimes \nu_2$$

which ends the proof. □

Remark 3.2. If $\mathcal{T}^{(1)}$ and $\mathcal{T}^{(2)}$ are weakly mixing and all ergodic joinings of $\mathcal{T}^{(1)}$ and $\mathcal{T}^{(2)}$ are weakly mixing then $\mathcal{J}(\mathcal{T}^{(1)}, \mathcal{T}^{(2)}) = \mathcal{J}(T_1^{(1)}, T_1^{(2)})$ and $\mathcal{J}^e(\mathcal{T}^{(1)}, \mathcal{T}^{(2)}) = \mathcal{J}^e(T_1^{(1)}, T_1^{(2)})$. Indeed, by Lemma 2.1 the only possibility to obtain a new ergodic joining is via suspension, every suspension however has a discrete spectrum factor.

Remark 3.3. If $\mathcal{J}_2(\mathcal{T}) = \mathcal{J}_2(T_1)$ then $\mathcal{T}$ and T_1 have the same invariant σ-algebras and $C(\mathcal{T}) = C(T_1)$.

Given an ergodic automorphism R on a standard probability Borel space $(Z, \mathcal{D}, \rho)$, we denote by $\mathcal{JF}(R)$ the family of all factors of all infinite ergodic self-joinings of R. Notice that

$$\begin{aligned} &\text{the class } \mathcal{JF}(R) \text{ is closed under} \\ &\text{(infinite) ergodic self-joinings and factors.} \end{aligned} \tag{14}$$

Therefore, for each ergodic automorphism S acting on a standard probability Borel space $(Y, \mathcal{C}, \nu)$ there exists a largest factor $\mathcal{A}_R(S) \subset \mathcal{C}$ of S such that $\mathcal{A}_R(S) \in \mathcal{JF}(R)$.[25] Since $S|_{W\mathcal{A}_R(S)}$ and $S|_{\mathcal{A}_R(S)}$ are isomorphic for each $W \in C(S)$, we obtain that

$$\mathcal{A}_R(S) \text{ is invariant under all } W \in C(S). \tag{15}$$

Now, fix a weakly mixing automorphism $R\colon (Z, \mathcal{D}, \rho) \to (Z, \mathcal{D}, \rho)$ and an ergodic automorphism $S\colon (Y, \mathcal{C}, \nu) \to (Y, \mathcal{C}, \nu)$. Then by (15), $\mathcal{A}_{R\times R}(S \times S)$ is a factor of the product action $(S^i \times S^j)_{(i,j)\in\mathbb{Z}^2}$. Recall a result on product $\mathbb{Z}^2$-actions:

Proposition 3.5. *[6] Let $\tilde{T}\colon (\tilde{X}, \tilde{\mathcal{B}}, \tilde{\mu}) \to (\tilde{X}, \tilde{\mathcal{B}}, \tilde{\mu})$ and $\tilde{S}\colon (\tilde{Y}, \tilde{\mathcal{C}}, \tilde{\nu}) \to (\tilde{Y}, \tilde{\mathcal{C}}, \tilde{\nu})$ be ergodic automorphisms of standard probability Borel spaces. Consider the corresponding $\mathbb{Z}^2$-action, i.e. the action of $(\tilde{T}^n \times \tilde{S}^m)_{n,m\in\mathbb{Z}^2}$ on $(\tilde{X} \times \tilde{Y}, \tilde{\mathcal{B}} \otimes$*

[25] By separability, the smallest σ-algebra including all factors of S from $\mathcal{JF}(R)$ is countably generated, whence it corresponds to a joining of at most countable number of factors.

$\tilde{\mathcal{C}}, \tilde{\mu}\otimes\tilde{\nu})$. *Suppose that* $\tilde{\mathcal{A}}$ *is a factor of the* $\mathbb{Z}^2$*-action. Let* $\overline{B}\otimes\overline{C}$ *be the smallest product factor containing* $\tilde{A}$.[26] *Then there exists a compact metric group* G *and two continuous 1-1 homomorphisms*

$$g \mapsto \tilde{R}_g \in C(\tilde{T}), \quad g \mapsto \tilde{R}'_g \in C(\tilde{S}),$$

such that

$$\tilde{A} = \{A \in \overline{B}\otimes\overline{C}\colon (\tilde{R}_g \times \tilde{R}'_g)A = A\}.$$

It follows from the above proposition and from (15) applied to $\mathcal{A}_{R\times R}(S\times S)$ that $\mathcal{A}_{R\times R}(S\times S)$ is a product-σ-algebra. Therefore

$$\mathcal{A}_{R\times R}(S\times S) = \mathcal{A}_R(S)\otimes\mathcal{A}_R(S) \tag{16}$$

as this is the largest product σ-algebra in $\mathcal{JF}(R\times R)$.

Moreover, using (14) and a result from [4] (see also [13]), we conclude that for any $\lambda \in \mathcal{J}^e(R,S)$ we have $\lambda = \big(\lambda|_{\mathcal{D}\otimes\mathcal{A}_R(S)}\big)\widehat{\ }$, i.e. λ is the relatively independent extension of $\lambda|_{\mathcal{D}\otimes\mathcal{A}_R(S)}$. In particular, using (16), for $\lambda \in \mathcal{J}^e(R, S\times S)$

$$\lambda = \big(\lambda|_{\mathcal{D}\otimes\mathcal{A}_R(S)\otimes\mathcal{A}_R(S)}\big)\widehat{\ }. \tag{17}$$

We will show now that in fact JP is an intrinsic property.

Proposition 3.6. *Let* R *be an ergodic automorphism such that in the class* $\mathcal{JF}(R)$ *there is some weakly mixing transformation. An automorphism* R *has the JP property if and only if for each weakly mixing* $S \in \mathcal{JF}(R)$ *and each* $\lambda \in J^e(R, S\times S)$ *we have either* $\lambda = (\lambda|_{Z\times Y_1})\otimes\nu$ *or* $\lambda = (\lambda|_{Z\times Y_2})\otimes\nu$.

Proof. Fix an arbitrary weakly mixing automorphism S and let $\lambda \in J^e(T, S\times S)$. Then (17) holds, equivalently, $\mathcal{D}$ and $\mathcal{C}\otimes\mathcal{C}$ are relatively independent over $\mathcal{A}_R(S)\otimes\mathcal{A}_R(S)$. Take $f \in L^\infty(Z,\rho)$, $g_i \in L^\infty(Y_i,\nu)$ for $i = 1,2$. Assume that $\int g_2 = 0$. We have

$$\begin{aligned}\int f\otimes g_1\otimes g_2\, d\lambda &= \int E^\lambda(f\otimes g_1\otimes g_2|\mathcal{A}_R(S)\otimes\mathcal{A}_R(S))\, d\lambda\\ &\overset{(17)}{=} \int E^\lambda(f)|\mathcal{A}_R(S)\otimes\mathcal{A}_R(S))E^\lambda(g_1\otimes g_2|\mathcal{A}_R(S)\otimes\mathcal{A}_R(S))\, d\lambda. \end{aligned}\tag{18}$$

To distinguish both σ-algebras $\mathcal{A}_R(S)$ we will write $\mathcal{A}^1_R(S)$ for $\mathcal{A}_R(S)\otimes\{\emptyset, Y\}$ and $\mathcal{A}^2_R(S)$ for $\{\emptyset, Y\}\otimes\mathcal{A}_R(S)$. By the assumption $\mathcal{D}\otimes\mathcal{A}^1_R(S) \perp \mathcal{A}^2_R(S)$ (up to a permutation of coordinates) with respect to λ, whence the last integral in (18) equals

$$\begin{aligned}&\int E^\lambda(f|\mathcal{A}^1_R(S))E^\nu(g_1|A^1_R(S))E^\nu(g_2|A^2_R(S))\, d\lambda\\ &\qquad\overset{(*)}{=} \left(\int E^\lambda(f|\mathcal{A}^1_R(S))E^\nu(g_1|\mathcal{A}^1_R(s))\, d\lambda\right)\cdot\int E^\nu(g_2|\mathcal{A}^2_R(S))\, d\lambda = 0,\end{aligned}$$

where $(*)$ follows by the independence of $\mathcal{A}^1_R(S)$ and $\mathcal{A}^2_R(S)$. This completes the proof. □

[26] The existence of such factor was also proved in [6].

Remark 3.4. Notice that if R is such that in the class $\mathcal{JF}(R)$ there are no weakly mixing automorphisms then in fact R is disjoint from all weakly mixing systems (see Theorem 4.8), which implies the JP property.

Remark 3.5. It was shown in [44] that there exists a non-weakly mixing automorphism, such that the class $\mathcal{JF}(R)$ includes some weakly mixing systems. This is done using Rokhlin extensions.[27]

Corollary 3.7. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be a measure-preserving flow whose all ergodic self-joinings are weakly mixing. Then T_1 has the JP property if and only if $\mathcal{T}$ has the JP property.*

Proof. This follows immediately by Remarks 3.2, 3.3 and Proposition 3.6. □

The following problem remains open:

Question 1. Does the JP property of a weakly mixing flow $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ imply the JP property for T_t, $t \neq 0$?

3.3 Isometric, distal and relatively weakly mixing extensions

3.3.1 Isometric extensions

Suppose that $T\colon (X, \mathcal{B}, \mu) \to (X, \mathcal{B}, \mu)$ is embeddable into some measurable flow $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$, i.e. $T = T_1$. Let $\{f'_n\colon n \in \mathbb{N}\} \subset L^2(X, \mu)$ be a dense subset and let $f_n := f'_n/\|f'_n\|_{L^2(X,\mu)}$ for $n \in \mathbb{N}$. Let $d_{\{f_n\}_{n\in\mathbb{N}}}$ be the metric defined as in (5), using $\{f_n\}_{n\in\mathbb{N}}$.

Lemma 3.8. *For any $\varepsilon > 0$ there exists $\delta > 0$ such that for all $t \in \mathbb{R}$ such that $|t| < \delta$ and $\xi \in \mathcal{J}_2(T)$*

$$\overline{d}_{\{f_n\}_{n\in\mathbb{N}}}(\xi \circ (T_t \times T_t), \xi) < \varepsilon/4.$$

Proof. Fix $\varepsilon > 0$. Let $N \in \mathbb{N}$ be such that $\sum_{i>N} 1/2^i < \varepsilon/32$. Let $\delta > 0$ be such that for $1 \leq i \leq N$ and $t \in \mathbb{R}$ such that $|t| < \delta$

$$\|f_i \circ T_t - f_i\|_{L^2(X,\mu)} < \varepsilon/8.$$

[27] Recall that the class of JP systems is closed under taking Rokhlin extensions.

Then for $t \in \mathbb{R}$ and $\xi \in \mathcal{J}_2(T)$ by Cauchy-Schwarz inequality we obtain

$$\begin{aligned}
d_{\{f_n\}_{n\in\mathbb{N}}}(\xi \circ (T_t \times T_t), \xi) &= \sum_{i,j\geq 1} \frac{1}{2^{i+j}} \left| \int_{X\times X} (f_i \circ T_t \otimes f_j \circ T_t - f_i \otimes f_j)\ d\xi \right| \\
&\leq \sum_{i,j\geq 1} \frac{1}{2^{i+j}} \left| \int_{X\times X} (f_i \circ T_t \otimes f_j \circ T_t - f_i \circ T_t \otimes f_j)\ d\xi \right| \\
&\quad + \sum_{i,j\geq 1} \frac{1}{2^{i+j}} \left| \int_{X\times X} (f_i \circ T_t \otimes f_j - f_i \otimes f_j)\ d\xi \right| \\
&\leq \sum_{i,j\geq 1} \frac{1}{2^{i+j}} \|f_j \circ T_t - f_j\|_{L^2(X,\mu)} + \sum_{i,j\geq 1} \frac{1}{2^{i+j}} \|f_i \circ T_t - f_i\|_{L^2(X,\mu)} \\
&= 2 \sum_{i\geq 1} \frac{1}{2^i} \|f_i \circ T_t - f_i\|_{L^2(X,\mu)} \\
&= 2 \left(\sum_{i=1}^{N} \frac{1}{2^i} \|f_i \circ T_t - f_i\|_{L^2(X,\mu)} + \sum_{i>N} \frac{1}{2^i} \|f_i \circ T_t - f_i\|_{L^2(X,\mu)} \right) \\
&\leq 2 \left(\sum_{i=1}^{N} \frac{1}{2^i} \cdot \varepsilon/8 + \varepsilon/16 \right) < \varepsilon/4,
\end{aligned}$$

which ends the proof. □

Let $A_\sigma = \{(m,p,q) \in \mathbb{N}^3 \colon m \geq 1, q \geq 0, 0 \leq p \leq 2^q - 1,\ p \text{ is odd}\}$ and $\sigma \colon A_\sigma \to \mathbb{N}$ be a bijection. For $i \geq 1$ let

$$g_i := f_m \circ T_{p/2^q} \text{ where } (m,p,q) = \sigma(i).$$

Then $\{g_n \colon n \in \mathbb{N}\} \subset L^2(X, \mathcal{B}, \mu)$ is clearly a dense subset as it includes $\{f_n \colon n \in \mathbb{N}\}$. Let $\overline{d}$ be the metric on $\mathcal{J}_2(T)$ defined as in (5), but using $\{g_n\}_{n\in\mathbb{N}}$ instead of $\{f_n\}_{n\in\mathbb{N}}$.

Lemma 3.9. *For any $\varepsilon > 0$ and any $q \in \mathbb{N}$ there exists $C = C(\varepsilon, q)$ such that for all $0 \leq p \leq 2^q - 1$ and all $\rho_1, \rho_2 \in \mathcal{J}_2(T)$*

$$\overline{d}\left(\rho_1 \circ (T_{p/2^q} \times T_{p/2^q}), \rho_2 \circ (T_{p/2^q} \times T_{p/2^q})\right) \leq C(\varepsilon, q) \cdot \overline{d}(\rho_1, \rho_2) + \varepsilon/2.$$

Proof. Fix $\varepsilon > 0$ and $q \in \mathbb{N}$. Let $0 \leq p \leq 2^q - 1$ and let $\rho_1, \rho_2 \in \mathcal{J}_2(T)$. Take $N \in \mathbb{N}$ such that

$$\sum_{i=1}^{N} \sum_{j>N} \frac{1}{2^{i+j}} + \sum_{i>N} \sum_{j=1}^{N} \frac{1}{2^{i+j}} < \varepsilon/2. \tag{19}$$

Let

$$a_{i,j} = \left| \int_{X\times X} g_i \otimes g_j\ d\rho_1 - \int_{X\times X} g_i \otimes g_j\ d\rho_2 \right|$$

for $i, j \geq 1$. Then

$$\overline{d}(\rho_1, \rho_2) = \sum_{i,j\geq 1} \frac{1}{2^{i+j}} a_{i,j}.$$

Let $\pi\colon \mathbb{N}\to\mathbb{N}$ be a bijection defined in the following way:

$$\pi(i)=j \iff \begin{cases}\sigma^{-1}(i)=(m,p_i,q_i),\\ \sigma^{-1}(j)=(m,p_j,q_j),\end{cases}$$
$$\text{for some } m\in\mathbb{N} \text{ and } q_i,q_j\in\mathbb{N},\ 0\le p_i\le 2^{q_i}-1,\ 0\le p_j\le 2^{q_j}-1$$
$$\text{such that } \frac{p_i}{2^{q_i}}+\frac{p}{2^q}-\frac{p_j}{2^{q_j}}\in\mathbb{Z}.$$

Permutation π is well-defined as the addition of $p/2^q$ mod 1 is a bijection of the set $\{p_i/2^{q_i}\colon (1,p_i,q_i)\in A_\sigma\}$. Recall also that ρ_1,ρ_2 are $T\times T$-invariant as elements of $\mathcal{J}_2(T)$. Therefore

$$\overline{d}(\rho_1\circ(T_{p/2^q}\times T_{p/2^q}),\rho_2\circ(T_{p/2^q}\times T_{p/2^q}))=\sum_{i,j\ge 1}\frac{1}{2^{i+j}}a_{\pi(i),\pi(j)},$$

By (19) we obtain

$$\begin{aligned}\sum_{i,j\ge1}\frac{1}{2^{i+j}}a_{\pi(i),\pi(j)} &\le \sum_{i,j=1}^{N}\frac{1}{2^{i+j}}a_{\pi(i),\pi(j)}+\varepsilon/2\\ &=\sum_{i,j=1}^{N}2^{\pi(i)-i+\pi(j)-j}\frac{1}{2^{\pi(i)+\pi(j)}}a_{\pi(i),\pi(j)}+\varepsilon/2\\ &\le \sum_{i,j=1}^{\max\{\pi(i)\colon 1\le i\le N\}}2^{2\max\{|\pi(i)-i|\colon 1\le i\le N\}}\frac{1}{2^{i+j}}a_{i,j}+\varepsilon/2\\ &\le 2^{2\max\{|\pi(i)-i|\colon 1\le i\le N\}}\sum_{i,j\ge1}\frac{1}{2^{i+j}}a_{i,j}+\varepsilon/2\\ &=C(\varepsilon,p,q)\cdot\overline{d}(\rho_1,\rho_2)+\varepsilon/2,\end{aligned}$$

where $C(\varepsilon,p,q)=2^{2\max\{|\pi(i)-i|\colon 1\le i\le N\}}$. To end the proof it suffices to put

$$C(\varepsilon,q):=\max\{C(\varepsilon,p,q)\colon 0\le p\le 2^q-1\}.$$

□

Lemma 3.10. *For any $\varepsilon>0$ there exists $\delta>0$ such that for any $\xi_1,\xi_2\in\mathcal{J}_2(T)$ if $\overline{d}(\xi_1,\xi_2)<\delta$ then for all $t\in\mathbb{R}$, $\overline{d}(\xi_1\circ(T_t\times T_t),\xi_2\circ(T_t\times T_t))<\varepsilon$. In particular, $\overline{d}\left(\int_0^1\xi_1\circ(T_t\times T_t)\,dt,\int_0^1\xi_2\circ(T_t\times T_t)\,dt\right)<\varepsilon$.*

Proof. Fix $\varepsilon>0$. By Lemma 3.8, we can find $\delta_0>0$ such that for $|t|<\delta_0$ and $\xi\in\mathcal{J}_2(T)$

$$\overline{d}(\xi\circ(T_t\times T_t),\xi)<\varepsilon/8. \tag{20}$$

Let $q\in\mathbb{N}$ be such that $1/2^q<\delta_0$ and let $C=C(\varepsilon,q_0)$ be such as in Lemma 3.9. For any $t\in\mathbb{R}$ find $0\le p\le 2^q-1$ such that $|t-p/2^q|<\delta_0$. Using (20) we obtain

$$\begin{aligned}&\overline{d}(\xi_1\circ(T_t\times T_t),\xi_2\circ(T_t\times T_t))\\ &\quad\le\overline{d}(\xi_1\circ(T_{p/2^q}\times T_{p/2^q}),\xi_2\circ(T_{p/2^q}\times T_{p/2^q}))+\varepsilon/4\\ &\quad\le C\cdot\overline{d}(\xi_1,\xi_2)+\varepsilon/4+\varepsilon/4=C\cdot\overline{d}(\xi_1,\xi_2)+\varepsilon/2,\end{aligned}$$

Let $0 < \delta < \frac{\varepsilon}{2 \cdot C}$. Then the condition $\overline{d}(\xi_1, \xi_2) < \delta$ implies

$$\overline{d}(\xi_1 \circ (T_t \times T_t), \xi_2 \circ (T_t \times T_t)) < \varepsilon.$$

It follows that

$$\overline{d}\left(\int_0^1 \xi_1 \circ (T_t \times T_t)\ dt, \int_0^1 \xi_2 \circ (T_t \times T_t)\ dt\right) \leq \int_0^1 \overline{d}(\xi_1 \circ (T_t \times T_t), \xi_2 \circ (T_t \times T_t))\ dt < \varepsilon,$$

which ends the proof. □

Proposition 3.11. *Let $\mathcal{T} = (T_t)_{t \in \mathbb{R}}$ be an ergodic flow on $(X, \mathcal{B}, \widetilde{\boldsymbol{\nu}})$ with a factor $\mathcal{A}$. Let $\underline{\boldsymbol{\nu}} = \widetilde{\boldsymbol{\nu}}|_{\mathcal{A}}$. Let $\widetilde{\boldsymbol{\nu}} = \int_0^1 \widetilde{\nu} \circ T_t\ dt$ and $\underline{\boldsymbol{\nu}} = \int_0^1 \nu \circ T_t|_{\mathcal{A}}\ dt$ for some $\widetilde{\nu} \in \mathcal{P}^e_{\mathbb{Z}}(T_1), \nu \in \mathcal{P}^e_{\mathbb{Z}}(T_1|_{\mathcal{A}})$.[28] Additionally assume that $\widetilde{\nu}|_{\mathcal{A}} = \nu$. Then the extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is isometric if and only if the extension $(\mathcal{T}, \widetilde{\boldsymbol{\nu}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\boldsymbol{\nu}})$ is isometric.*

Proof. Suppose that the extension $(\mathcal{T}, \widetilde{\boldsymbol{\nu}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\boldsymbol{\nu}})$ is isometric. Let $\overline{\mathcal{T}} = (\overline{T}_t)_{t \in \mathbb{R}} \colon (\overline{X}, \overline{\mathcal{B}}, \overline{\boldsymbol{\nu}}) \to (\overline{X}, \overline{\mathcal{B}}, \overline{\boldsymbol{\nu}})$ be an ergodic extension of $\mathcal{T}$ such that $(\overline{\mathcal{T}}, \overline{\boldsymbol{\nu}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\boldsymbol{\nu}})$ is a compact group extension. By Lemma 2.1 there exists $\overline{\nu} \in \mathcal{P}^e_{\mathbb{Z}}(\overline{T}_1)$ such that $\overline{\boldsymbol{\nu}} = \int_0^1 \overline{\nu} \circ \overline{T}_t\ dt$ (it may happen that $\overline{\boldsymbol{\nu}} = \overline{\nu}$). By the uniqueness of the ergodic decomposition and by Lemma 2.2, there exist $k, l, m \in \mathbb{N} \cup \{\infty\}$ such that

$$\overline{\boldsymbol{\nu}} = k \int_0^{1/k} \overline{\nu} \circ \overline{T}_t\ dt, \quad \widetilde{\boldsymbol{\nu}} = kl \int_0^{1/kl} \widetilde{\nu} \circ T_t\ dt, \quad \underline{\boldsymbol{\nu}} = klm \int_0^{1/klm} \nu \circ T_t|_{\mathcal{A}}\ dt.^{29}$$

Without loss of generality we may assume that $\overline{\nu}|_{\mathcal{B}} = \widetilde{\nu}$. Since $(\overline{\mathcal{T}}, \overline{\boldsymbol{\nu}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\boldsymbol{\nu}})$ is a compact group extension, $(\overline{T}_1, \overline{\boldsymbol{\nu}}) \to (T_1|_{\mathcal{A}}, \underline{\boldsymbol{\nu}})$ is also a compact group extension. Notice that in the decomposition $\overline{\boldsymbol{\nu}} = k \int_0^{1/k} \overline{\nu} \circ \overline{T}_t\ dt$ the only measures which project down to ν are measures of the form $\overline{\nu} \circ T_{i/lm}$ for $0 \leq i \leq lm - 1$. Therefore also

$$\left(\overline{T}_1, \frac{1}{lm} \sum_{i=0}^{lm-1} \overline{\nu} \circ T_{i/lm}\right) \to (T_1|_{\mathcal{A}}, \nu)$$

is a compact group extension (if $lm = \infty$, we interpret $\frac{1}{lm} \sum_{i=0}^{lm-1} \overline{\nu} \circ T_{i/lm}$ as $\overline{\boldsymbol{\nu}}$). It follows that also $(\overline{T}_1, \overline{\nu}) \to (T_1, \nu)$ is a compact group extension. Thus $(T_1, \widetilde{\nu}) \to (T_1, \nu)$ is isometric which ends the first part of the proof.

Assume now that the extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is isometric. As in the first part of the proof, there exist $k, l \in \mathbb{N} \cup \{\infty\}$ such that

$$\widetilde{\boldsymbol{\nu}} = k \int_0^{1/k} \widetilde{\nu} \circ T_t\ dt, \ \underline{\boldsymbol{\nu}} = kl \int_0^{1/kl} \nu \circ T_t|_{\mathcal{A}}\ dt \tag{21}$$

[28] It may happen that $\underline{\boldsymbol{\nu}} = \nu$ or $\widetilde{\boldsymbol{\nu}} = \widetilde{\nu}$. The existence of ν and $\widetilde{\nu}$ follows from Lemma 2.1.

[29] We interpret $\int_0^{1/\infty} \underline{\boldsymbol{\lambda}} \circ S_t\ dt$ as $\underline{\boldsymbol{\lambda}}$ for any measure $\underline{\boldsymbol{\lambda}}$ and any flow $(S_t)_{t \in \mathbb{R}}$ such that this formula makes sense.

are the corresponding ergodic decompositions (see footnotes 28 and 29). By Proposition 2.8 the extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is T-compact. Fix $\varepsilon > 0$. Let $0 < \delta < \varepsilon$ be as in Lemma 3.8. Let $N \in \mathbb{N}$ come from the definition of T-compactness for $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ with ε replaced with δ. Take $\underline{\boldsymbol{\lambda}} \in J_N^e((T, \underline{\widetilde{\boldsymbol{\nu}}}); \mathcal{A})$. Then $\underline{\boldsymbol{\lambda}} \in \mathcal{P}_{\mathbb{Z}}(T_1^{\times N})$ and there are two possibilities:

(i) $\underline{\boldsymbol{\lambda}} \in \mathcal{P}_{\mathbb{Z}}^e(T_1^{\times N})$,

(ii) $\underline{\boldsymbol{\lambda}} \notin \mathcal{P}_{\mathbb{Z}}^e(T_1^{\times N})$.

Case (i) may occur only if $k = \infty$, i.e. $\underline{\widetilde{\boldsymbol{\nu}}} = \widetilde{\nu}$, $\underline{\boldsymbol{\nu}} = \nu$ (otherwise the marginals of $\underline{\boldsymbol{\lambda}}$ are not ergodic for T_1, so $\underline{\boldsymbol{\lambda}}$ is not ergodic for $T_1^{\times N}$). Then by Proposition 2.8 there exist $1 \leq i < j \leq N$ such that

$$\overline{d}(\underline{\boldsymbol{\lambda}}|_{X_i \times X_j}, \Delta_{X_i \times X_j}(\underline{\widetilde{\boldsymbol{\nu}}})) < \delta < \varepsilon.$$

We will cover now case (ii). To fix our attention, we will assume that $k, l \in \mathbb{N}$, i.e. $\underline{\widetilde{\boldsymbol{\nu}}} \neq \widetilde{\nu}$ and $\underline{\boldsymbol{\nu}} \neq \nu$ (in other cases the proof is similar). By Lemma 2.1 there exists $\lambda \in \mathcal{P}_{\mathbb{Z}}^e(T_1^{\times N})$ such that

$$\underline{\boldsymbol{\lambda}} = \int_0^1 \lambda \circ T_t^{\times N} \, dt. \tag{22}$$

We will show that

$$\text{such a measure } \lambda \text{ can be chosen from } \mathcal{J}_N^e((T_1, \widetilde{\nu}); \mathcal{A}). \tag{23}$$

In what follows we will use the fact that ergodic decompositions are unique. We have the following:

$$\begin{aligned} \underline{\boldsymbol{\lambda}}|_{\mathcal{A}^{\otimes N}} &= \int_0^1 \lambda|_{\mathcal{A}^{\otimes N}} \circ (T_t|_{\mathcal{A}})^{\times N} \, dt, \\ \underline{\boldsymbol{\lambda}}|_{\mathcal{A}^{\otimes N}} &= \Delta_{\mathcal{A}^{\otimes N}}(\underline{\boldsymbol{\nu}}) = kl \int_0^{1/kl} \Delta_{\mathcal{A}^{\otimes N}}(\nu) \circ (T_t|_{\mathcal{A}})^{\times N} \, dt \end{aligned} \tag{24}$$

(the latter equality follows from the second formula in (21)). Therefore we may assume without loss of generality (changing $\lambda \in \mathcal{P}_{\mathbb{Z}}^e(T_1^{\times N})$ if necessary) that

$$\lambda|_{\mathcal{A}^{\otimes N}} = \Delta_{\mathcal{A}^{\otimes N}}(\nu) \tag{25}$$

and $\underline{\boldsymbol{\lambda}}|_{\mathcal{A}^{\otimes N}} = kl \int_0^{1/kl} \lambda|_{\mathcal{A}^{\otimes N}} \circ (T_t|_{\mathcal{A}})^{\times N} \, dt$. Moreover, for $1 \leq i \leq N$ we have

$$\begin{aligned} \underline{\boldsymbol{\lambda}}|_{X_i} &= k \int_0^{1/k} \lambda|_{X_i} \circ T_t \, dt, \\ \underline{\boldsymbol{\lambda}}|_{X_i} &= \underline{\widetilde{\boldsymbol{\nu}}} = k \int_0^{1/k} \widetilde{\nu} \circ T_t \, dt, \end{aligned}$$

whence there exists $t_i \in [0, 1/k)$ such that

$$\lambda|_{X_i} = \widetilde{\nu} \circ T_{t_i}. \tag{26}$$

It follows that

$$(\lambda|_{X_i})|_{\mathcal{A}} = \widetilde{\nu}|_{\mathcal{A}} \circ T_{t_i}|_{\mathcal{A}} = \nu \circ (T_{t_i}|_{\mathcal{A}}).$$

On the other hand by (25)

$$(\lambda|_{X_i})\,|_{\mathcal{A}} = \nu.$$

Hence $t_i = 0$ and (23) follows from (25) and (26). By Proposition 2.8 there exist $1 \le i < j \le N$ such that

$$\overline{d}(\lambda|_{X_i\times X_j}, \Delta_{X_i\times X_j}(\widetilde{\nu})) < \delta. \tag{27}$$

By (22) we have

$$\underline{\boldsymbol{\lambda}}|_{X_i\times X_j} = \int_0^1 \lambda|_{X_i\times X_j} \circ (T_t \times T_t)\; dt. \tag{28}$$

Moreover,

$$\Delta_{X_i\times X_j}(\underline{\widetilde{\boldsymbol{\nu}}}) = \int_0^1 \Delta_{X_i\times X_j}(\widetilde{\nu}) \circ (T_t \times T_t)\; dt \tag{29}$$

It follows from the choice of δ, from (27), (28) and (29) and from Lemma 3.10 that

$$\overline{d}(\underline{\boldsymbol{\lambda}}|_{X_i\times X_j}, \Delta_{X_i\times X_j}(\underline{\widetilde{\boldsymbol{\nu}}})) < \varepsilon.$$

The claim follows from Proposition 2.8. □

Proposition 3.12. *Let $l \in \mathbb{N}$ and let T be an ergodic automorphism of $(X, \mathcal{B}, \underline{\widetilde{\boldsymbol{\nu}}})$ with a factor $\mathcal{A}$. Let $\underline{\boldsymbol{\nu}} = \underline{\widetilde{\boldsymbol{\nu}}}|_{\mathcal{A}}$. Let $\underline{\widetilde{\boldsymbol{\nu}}} = \frac{1}{l}\sum_{i=1}^{l-1} \widetilde{\nu}\circ T^i$ and $\underline{\boldsymbol{\nu}} = \frac{1}{l}\sum_{i=0}^{l-1} \nu\circ T^i|_{\mathcal{A}}$ for some $\widetilde{\nu} \in \mathcal{P}^e_{\mathbb{Z}}(T^l), \nu \in \mathcal{P}^e_{\mathbb{Z}}(T^l|_{\mathcal{A}})$. Additionally assume that $\widetilde{\nu}|_{\mathcal{A}} = \nu$. Then the extension $(T^l, \widetilde{\nu}) \to (T^l|_{\mathcal{A}}, \nu)$ is isometric if and only if the extension $(T, \underline{\widetilde{\boldsymbol{\nu}}}) \to (T|_{\mathcal{A}}, \underline{\boldsymbol{\nu}})$ is isometric.*

Proof. The proof goes along the same lines as the proof of Proposition 3.11. □

3.3.2 Distal extensions and Furstenberg-Zimmer decomposition

Distal extensions via isometric extensions

Proposition 3.13. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be an ergodic flow on $(X, \mathcal{B}, \underline{\widetilde{\boldsymbol{\nu}}})$ with a factor $\mathcal{A}$. Let $\underline{\boldsymbol{\nu}} = \underline{\widetilde{\boldsymbol{\nu}}}|_{\mathcal{A}}$. Let $\underline{\widetilde{\boldsymbol{\nu}}} = \int_0^1 \widetilde{\nu}\circ T_t\; dt$ and $\underline{\boldsymbol{\nu}} = \int_0^1 \nu\circ T_t\; dt$ for some $\widetilde{\nu} \in \mathcal{P}^e_{\mathbb{Z}}(T_1), \nu \in \mathcal{P}^e_{\mathbb{Z}}(T_1|_{\mathcal{A}})$.[30] Additionally assume that $\widetilde{\nu}|_{\mathcal{A}} = \nu$. Then the extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is distal if and only if the extension $(\mathcal{T}, \underline{\widetilde{\boldsymbol{\nu}}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\boldsymbol{\nu}})$ is distal.*

Proof. Suppose that the extension $(\mathcal{T}, \underline{\widetilde{\boldsymbol{\nu}}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\boldsymbol{\nu}})$ is distal. Since any factor of $\mathcal{T}$ is a factor of T_1, it clearly follows from the definition of a distal extension and Proposition 3.11 that also the extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is distal.

Suppose now that the extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is distal. Let $(\mathcal{C}_\alpha)_{\alpha\le\beta}$ be a transfinite sequence of factors of T_1 such that $\mathcal{C}_0 = \mathcal{A}$, $\mathcal{C}_\beta = \mathcal{B}$, the extension $\mathcal{C}_{\alpha+1} \to \mathcal{C}_\alpha$ for T_1 is isometric and for α being limit ordinals, $\mathcal{C}_\alpha$ is the inverse limit of the preceding factors. We may assume that for each α the factor $\mathcal{C}_{\alpha+1}$ is the largest extension of $\mathcal{C}_\alpha$ inside $\mathcal{B}$ such that $\mathcal{C}_{\alpha+1} \to \mathcal{C}_\alpha$ is isometric. In particular, for each α, $\mathcal{C}_\alpha$ is $\mathcal{T}$-invariant, i.e. it is a factor of $\mathcal{T}$. Indeed, for α which is not a limit ordinal, $T_1|_{\mathcal{C}_\alpha}$ is isomorphic to $T_1|_{T_t\mathcal{C}_\alpha}$, both $\mathcal{C}_\alpha \to \mathcal{C}_{\alpha-1}$, $T_t\mathcal{C}_\alpha \to \mathcal{C}_{\alpha-1}$ are isometric, whence $\mathcal{C}_\alpha \vee T_t\mathcal{C}_\alpha \to \mathcal{C}_{\alpha-1}$ is also isometric, which

[30] See footnote 29.

follows e.g. from (8).[31] For α which is a limit ordinal it suffices to notice that the property of being $\mathcal{T}$-invariant is preserved under taking inverse limits. This ends the proof by Proposition 3.11. □

The same proof works in the co-finite case:

Proposition 3.14. *Let T be an ergodic automorphism acting on $(X, \mathcal{B}, \widetilde{\underline{\nu}})$ with a factor $\mathcal{A}$. Fix $l \in \mathbb{N}$. Let $\underline{\nu} = \widetilde{\underline{\nu}}|_{\mathcal{A}}$. Let $\widetilde{\underline{\nu}} = \frac{1}{l}\sum_{i=0}^{l-1} \widetilde{\nu} \circ T_i$ and $\underline{\nu} = \frac{1}{l}\sum_{i=0}^{l-1} \nu \circ T_i|_{\mathcal{A}}$ for some $\widetilde{\nu} \in \mathcal{P}^e_{\mathbb{Z}}(T^l), \nu \in \mathcal{P}^e_{\mathbb{Z}}(T^l|_{\mathcal{A}})$. Additionally assume that $\widetilde{\nu}|_{\mathcal{A}} = \nu$. Then the extension $(T^l, \widetilde{\nu}) \to (T^l|_{\mathcal{A}}, \nu)$ is distal if and only if the extension $(T, \widetilde{\underline{\nu}}) \to (T|_{\mathcal{A}}, \underline{\nu})$ is distal.*

Furstenberg-Zimmer decomposition Using the techniques from the proof of Proposition 3.13 and reasoning by contradiction one can show more:

Corollary 3.15. *Under the assumptions of Proposition 3.13 the following statements are true and equivalent:*

(i) The extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is distal if and only if the extension $(\mathcal{T}, \widetilde{\underline{\nu}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\nu})$ is distal.

(ii) The extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is relatively weakly mixing if and only if the extension $(\mathcal{T}, \widetilde{\underline{\nu}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\nu})$ is relatively weakly mixing.

(iii) The Furstenberg-Zimmer decomposition for $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is the same as for $(\mathcal{T}, \widetilde{\underline{\nu}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\nu})$, i.e. there exists $\mathcal{A} \subset \mathcal{C} \subset \mathcal{B}$ which is a factor of $\mathcal{T}$ such that the extension $\mathcal{B} \to \mathcal{C}$ is relatively weakly mixing both for $(\mathcal{T}, \widetilde{\underline{\nu}})$ and for $(T_1, \widetilde{\nu})$ and the extension $\mathcal{C} \to \mathcal{A}$ is distal both for $(\mathcal{T}, \widetilde{\underline{\nu}}|_{\mathcal{C}})$ and for $(T_1, \widetilde{\nu}|_{\mathcal{C}})$.

In the co-finite case we have the following:

Corollary 3.16. *Under the assumptions of Proposition 3.14 the following statements are true and equivalent:*

(i) The extension $(T^l, \widetilde{\nu}) \to (T^l|_{\mathcal{A}}, \nu)$ is distal if and only if the extension $(T, \widetilde{\underline{\nu}}) \to (T|_{\mathcal{A}}, \underline{\nu})$ is distal.

(ii) The extension $(T^l, \widetilde{\nu}) \to (T^l|_{\mathcal{A}}, \nu)$ is relatively weakly mixing if and only if the extension $(T, \widetilde{\underline{\nu}}) \to (T|_{\mathcal{A}}, \underline{\nu})$ is relatively weakly mixing.

(iii) Then the Furstenberg-Zimmer decomposition for $(T^l, \widetilde{\nu}) \to (T^l|_{\mathcal{A}}, \nu)$ is the same as for $(T, \widetilde{\underline{\nu}}) \to (T|_{\mathcal{A}}, \underline{\nu})$, i.e. there exists $\mathcal{A} \subset \mathcal{C} \subset \mathcal{B}$ which is a factor of T_1 such that the extension $\mathcal{B} \to \mathcal{C}$ is relatively weakly mixing both for $(T, \widetilde{\underline{\nu}})$ and for $(T^l, \widetilde{\nu})$ and the extension $\mathcal{C} \to \mathcal{A}$ is distal both for $(T, \widetilde{\underline{\nu}}|_{\mathcal{C}})$ and for $(T^l, \widetilde{\nu}|_{\mathcal{C}})$.

[31] $\mathcal{C}_\alpha \vee T_t\mathcal{C}_\alpha$ stands for the σ-algebra generated by $\mathcal{C}_\alpha$ and $T_t\mathcal{C}_\alpha$.

Distal extensions - independent approach Let G be a locally compact, second countable abelian group and let H be its co-compact subgroup. Suppose G acts ergodically on $(X, \mathcal{B}, \mu)$. Let $\mathcal{A} \subset \mathcal{B}$ be a factor of G such that the subaction of H restricted to $(X, \mathcal{A}, \mu|_{\mathcal{A}})$ is ergodic. There are two possibilities:

(A_G) the H-sub-action is ergodic with respect to μ,

(B_G) the H-sub-action is not ergodic with respect to μ.

We will be interested in $G \in \{\mathbb{R}, \mathbb{Z}\}$ and $H = l\mathbb{Z}$ for $l \in \mathbb{N}$. In order to give an alternative proof of Proposition 3.13 and Proposition 3.14 in case $\underline{\boldsymbol{\nu}} = \nu$[32] we take the following path:

$$(A_{\mathbb{Z}}) \to [(B_{\mathbb{R}}) \text{ and } (B_{\mathbb{Z}})] \to (A_{\mathbb{R}}).$$

Notice that

- case $(A_{\mathbb{Z}})$ corresponds to Proposition 3.14 in case $\underline{\widetilde{\boldsymbol{\nu}}} = \widetilde{\nu}$,
- case $(B_{\mathbb{R}})$ corresponds to Proposition 3.13 in case $\underline{\widetilde{\boldsymbol{\nu}}} \neq \widetilde{\nu}$,
- case $(B_{\mathbb{Z}})$ corresponds to Proposition 3.14 in case $\underline{\widetilde{\boldsymbol{\nu}}} \neq \widetilde{\nu}$,
- case $(A_{\mathbb{R}})$ corresponds to Proposition 3.13 in case $\underline{\widetilde{\boldsymbol{\nu}}} = \widetilde{\nu}$,

Case $(A_{\mathbb{Z}})$

Proof of Proposition 3.14 in case $\underline{\widetilde{\boldsymbol{\nu}}} = \widetilde{\nu}$. We will use Remark 2.6 on page 16 and use the notion of a separating sieve instead of the notion of relative distality. It is clear that every separating sieve for $(T, \widetilde{\nu}) \to (T|_{\mathcal{A}}, \nu)$ is a separating sieve for $(T^l, \widetilde{\nu}) \to (T^l|_{\mathcal{A}}, \nu)$. Let $A_1 \supset A_2 \supset \dots$ be a separating sieve for $(T^l, \widetilde{\nu}) \to (T^l|_{\mathcal{A}}, \nu)$ and let $X_0 \subset X$ be the set of full measure taken from the definition of this separating sieve. Let π be the map yielding the factor σ-algebra $\mathcal{A}$. Define $\widetilde{X}$ as the intersection $\cap_{k\in\mathbb{Z}} T^k X_0$. Take $\widetilde{x}, \widetilde{y} \in \widetilde{X}$ with $\pi(\widetilde{x}) = \pi(\widetilde{y})$ and suppose that for all $n \in \mathbb{N}$ there exists $k_n \in \mathbb{Z}$ such that $T^{k_n}\widetilde{x}, T^{k_n}\widetilde{y} \in A_n$. Let $l_n \in \{0, 1, \dots, l-1\}$ be such that $k_n - l_n \in l\mathbb{Z}$. Then

$$T^{k_n - l_n} T^{l_n} \widetilde{x}, T^{k_n - l_n} T^{l_n} \widetilde{y} \in A_n.$$

Since all l_n belong to the finite set $\{0, 1, \dots, l-1\}$ therefore along some subsequence $(n_k)_{k\in\mathbb{N}}$ we have l_{n_k} constant. Moreover $\pi(T^{l_{n_k}}\widetilde{x}) = \pi(T^{l_{n_k}}\widetilde{y})$ (π is the factoring map) and $T^{l_{n_k}}\widetilde{x}, T^{l_{n_k}}\widetilde{y} \in \widetilde{X}$. Our claim follows by the assumption that $A_1 \supset A_2 \supset \dots$ is a separating sieve for $(T^l, \widetilde{\nu}) \to (T^l|_{\mathcal{A}}, \nu)$. □

Cases $(B_{\mathbb{R}})$ and $(B_{\mathbb{Z}})$

Proof of Proposition 3.13 in case $\underline{\widetilde{\boldsymbol{\nu}}} \neq \widetilde{\nu}$ *(and* $\underline{\boldsymbol{\nu}} = \nu$*).* By Lemma 2.2 the ergodic decomposition of $\underline{\widetilde{\boldsymbol{\nu}}}$ for T_1 is of the form

$$\underline{\widetilde{\boldsymbol{\nu}}} = k \int_0^{1/k} \widetilde{\nu} \circ T_t \, dt$$

[32] This is the situation we will later observe in applications.

for some $k \in \mathbb{N}$ (the measures $\widetilde{\nu} \circ T_t$ are pairwise orthogonal for $t \in [0,1)$ and $\widetilde{\nu}$ is $T_{1/k}$-invariant). Moreover, $\widetilde{\nu}$ is ergodic for $T_{1/k}$ as it is ergodic for T_1. By Proposition 3.14 in case $(A_{\mathbb{Z}})$ proved above, the extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is distal if and only if the extension $(T_{1/k}, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is distal. Therefore we may assume without loss of generality that $k = 1$ and

$$\text{the measures } \widetilde{\nu} \circ T_t \text{ are mutually singular for } t \in [0,1). \tag{30}$$

Moreover, using Remark 2.3 on page 11, we may also assume that

$$\begin{aligned}(X, \mathcal{B}, \underline{\widetilde{\boldsymbol{\nu}}}) = (X, \mathcal{B}, \widetilde{\nu}) \times [0,1],\\ \text{with the identification of } (x,1) \text{ with } (T_1 x, 0)\\ \text{and } T_t(x,s) = (x, s+t) \text{ for } s \in [0,1].\end{aligned} \tag{31}$$

Suppose that the extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is distal. We will show that also the extension $(\mathcal{T}, \underline{\widetilde{\boldsymbol{\nu}}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\boldsymbol{\nu}})$ is distal. Since every ergodic factor of a non-ergodic group action is a factor of its almost every ergodic component we have $\pi(x,t) = \pi(x,s)$ for $0 \le t, s < 1$, where π is the factoring map. Let $X \times \{0\} \supset A_1 \supset A_2 \supset \dots$ be a separating sieve for $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$. Let X_0 be the set of full measure taken from the definition of the separating sieve for this extension. Take $\widetilde{x} = (x,r), \widetilde{y} = (y,s) \in X_0 \times [0,1]$. Suppose that for all $n \in \mathbb{N}$ there exists $t_n \in \mathbb{R}$ such that

$$T_{t_n}\widetilde{x}, T_{t_n}\widetilde{y} \in B_n := A_n \times [0, 1/n].$$

Then

$$T_{t_n - r_n}\widetilde{x} = T_{t_n - r_n + r}(x,0), T_{t_n - s_n}\widetilde{y} = T_{t_n - s_n + s}(y,0) \in A_n \times \{0\}$$

for some $r_n, s_n \in [0, 1/n]$. It follows by (30) that $t_n - r_n + r, t_n - s_n + s \in \mathbb{Z}$, whence $r_n - s_n + s - r \in \mathbb{Z}$. Since $r_n - s_n \in [-1/n, 1/n]$ and $s - r \in (-1, 1)$ therefore there exists $n_0 \in \mathbb{N}$ such that for $n \ge n_0$ we have $r_n - s_n + s - r = 0$, i.e. $r_n - s_n = r - s = 0$. Moreover, since $X \times \{0\} \supset A_1 \supset A_2 \supset \dots$ is a separating sieve for $(T_1, \widetilde{\nu}) \to (T_1, \nu)$, $x = y$. Hence $\widetilde{x} = \widetilde{y}$ and $B_1 \supset B_2 \supset \dots$ is a separating sieve for $(\mathcal{T}, \underline{\widetilde{\boldsymbol{\nu}}}) \to (\mathcal{T}, \underline{\boldsymbol{\nu}})$, which ends the first part of the proof.

Suppose now that the extension $(\mathcal{T}, \underline{\widetilde{\boldsymbol{\nu}}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\boldsymbol{\nu}})$ is distal. We will show that also the extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is distal. By the assumption

$$\text{there exists a separating sieve for } (\mathcal{T}, \underline{\widetilde{\boldsymbol{\nu}}}) \to (\mathcal{T}|_{\mathcal{A}}, \underline{\boldsymbol{\nu}}),$$

i.e. there exists a set $\widetilde{X}_0 \subset X$ with $\underline{\widetilde{\boldsymbol{\nu}}}(\widetilde{X}_0) = 1$ and a sequence of sets $(A_n)_{n\in\mathbb{N}}$ with $\underline{\widetilde{\boldsymbol{\nu}}}(A_n) > 0$ and $\underline{\widetilde{\boldsymbol{\nu}}}(A_n) \to 0$ such that whenever $x, y \in \widetilde{X}_0$ are such that $\pi(x) = \pi(y)$ and for every $n \in \mathbb{N}$ there exists $t_n \in \mathbb{R}$ with $T_{t_n}x, T_{t_n}y \in A_n$ then $x = y$.

We will use $(A_n)_{n\in\mathbb{N}}$ to find a separating sieve for $(T_1, \widetilde{\nu}) \to (T_1, \nu)$. Without loss of generality we may assume that $\widetilde{X}_0$ is invariant under the flow $(T_t)_{t\in\mathbb{R}}$, whence $\widetilde{\nu}(X_0) = 1$. Let

$$B_n := T_{t_n} A_n, \text{ where } t_n \in [0,1) \text{ is such that } \widetilde{\nu}(B_n) > 0.$$

We claim that $\widetilde{X}_0$ and $(B_n)_{n\in\mathbb{N}}$ satisfy the assumptions of Lemma 2.7. Indeed, let $x, y \in \widetilde{X}_0$ be such that $\pi(x) = \pi(y)$ and for all $n \in \mathbb{N}$ there exists $k_n \in \mathbb{Z}$ with $T_{k_n}x, T_{k_n}y \in B_n$. Then

$$T_{k_n - t_n}x, T_{k_n - t_n}y \in A_n,$$

which implies that $x = y$. To end the proof it suffices to notice that the measure $\widetilde{\nu}$ is non-atomic ($\underline{\widetilde{\boldsymbol{\nu}}} = \int_0^1 \widetilde{\nu} \circ T_t \, dt$ and $\underline{\widetilde{\boldsymbol{\nu}}}$ is finite) and use Lemma 2.7. □

In case $(B_{\mathbb{Z}})$ the proofs are the same as in case $(B_{\mathbb{R}})$.

Case $(A_{\mathbb{R}})$

Proof of Proposition 3.13 in case $\underline{\widetilde{\boldsymbol{\nu}}} = \widetilde{\nu}$. It is clear that every separating sieve for $(\mathcal{T}, \widetilde{\nu}) \to (\mathcal{T}|_{\mathcal{A}}, \nu)$ is a separating sieve for $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$. Using Remark 2.6 on page 16 we obtain the first part of the claim.

Suppose now that the extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is isometric. By the proof of Proposition 2.8 (see [47]) it follows that there exists $\kappa \in \mathcal{J}^e_\infty((T_1, \widetilde{\nu}); \mathcal{A})$ such that

$$(T_1^{\times\infty}, \kappa) \to (T_1|_{\mathcal{A}}, \nu) \text{ is a compact group extension.}$$

On the space $X^{\times\infty}$ we have a natural action of $\mathcal{T}^{\times\infty}$. It is however not necessarily true that the measure κ is $\mathcal{T}^{\times\infty}$-invariant. Hence we consider two cases:

(i) κ is $\mathcal{T}^{\times\infty}$-invariant,

(ii) κ is not $\mathcal{T}^{\times\infty}$-invariant.

In case (i) we have the following two possibilities:

(i1) $\mathcal{J}^e_2((\mathcal{T}^{\times\infty}, \kappa); \mathcal{A}) \subset \mathcal{J}^e_2((T_1^{\times\infty}, \kappa); \mathcal{A})$,

(i2) $\mathcal{J}^e_2((\mathcal{T}^{\times\infty}, \kappa); \mathcal{A}) \not\subset \mathcal{J}^e_2((T_1^{\times\infty}, \kappa); \mathcal{A})$.

In case (i1) by Proposition 2.3 and Proposition 2.4 we obtain that $(\mathcal{T}^{\times\infty}, \kappa) \to (\mathcal{T}|_{\mathcal{A}}, \nu)$ is a compact group extension. Therefore $(\mathcal{T}, \widetilde{\nu}) \to (\mathcal{T}|_{\mathcal{A}}, \nu)$ is an isometric extension. In case (i2) take

$$\underline{\boldsymbol{\xi}} \in \mathcal{J}^e_2((\mathcal{T}^{\times\infty}, \kappa); \mathcal{A}) \setminus \mathcal{J}^e_2((T_1^{\times\infty}, \kappa); \mathcal{A}).$$

Then

$$\underline{\boldsymbol{\xi}} = \int_0^1 \xi \circ \left(T_t^{\times\infty}\right)^{\times 2} \, dt, \text{ where } \xi \in \mathcal{J}^e_2((T_1^{\times\infty}, \kappa); \mathcal{A}).$$

By Proposition 2.3 the measure ξ is a graph self-joining, i.e.

$$(T_1^{\times\infty} \times T_1^{\times\infty}, \xi) \simeq (T_1^{\times\infty}, \kappa).$$

Therefore

$$(T_1^{\times\infty} \times T_1^{\times\infty}, \xi) \to (T_1, \nu) \text{ is a compact group extension,}$$

i.e. it is distal. It follows from Proposition 3.13 in case $(B_{\mathbb{R}})$ proved above, that

$$(\mathcal{T}^{\times\infty} \times \mathcal{T}^{\times\infty}, \underline{\boldsymbol{\xi}}) \to (\mathcal{T}|_{\mathcal{A}}, \nu) \text{ is also distal.}$$

Hence $(\mathcal{T}, \widetilde{\nu}) \to (\mathcal{T}|_{\mathcal{A}}, \nu)$ is also distal.

In case (ii) consider

$$\underline{\boldsymbol{\kappa}} := \int_0^1 \kappa \circ T_t^{\times\infty} \, dt.$$

By Proposition 3.13 in case $(B_{\mathbb{R}})$ it follows that $(\mathcal{T}^{\times\infty}, \underline{\boldsymbol{\kappa}}) \to (\mathcal{T}|_{\mathcal{A}}, \nu)$ is distal, whence $(\mathcal{T}, \widetilde{\nu}) \to (\mathcal{T}|_{\mathcal{A}}, \nu)$ is also a distal extension.

If the extension $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ is distal (and not isometric), it suffices to consider the maximal isometric extensions contributing to $(T_1, \widetilde{\nu}) \to (T_1|_{\mathcal{A}}, \nu)$ and use a similar "maximizing" procedure as in the proof of Proposition 3.13 (page 27). □

3.4 Quasi- and distal simplicity

In this section we prove counterparts of Proposition 3.1 and Proposition 3.4 for the 2-QS and 2-DS properties.

3.4.1 Quasi-simplicity – general case

Proposition 3.17. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be a measure-preserving flow on $(X, \mathcal{B}, \mu)$ with T_1 ergodic. Then $\mathcal{T}$ is 2-QS if and only if T_1 is 2-QS.*

Proof. Suppose that $\mathcal{T}$ is 2-QS and let $\lambda \in \mathcal{J}_2^e(T_1)$, $\lambda \neq \mu \otimes \mu$. There are two possibilities:

(i) $\lambda = \lambda \circ (T_t \times T_t)$ for all $t \in \mathbb{R}$,

(ii) $\lambda \perp \lambda \circ (T_t \times T_t)$ for some $t \in (0, 1)$.

In case (i) apply Proposition 3.11 to $(\mathcal{T} \times \mathcal{T}, \lambda)$, $(T_1 \times T_1, \lambda)$ and the coordinate factors. In case (ii) consider

$$\underline{\boldsymbol{\lambda}} := \int_0^1 \lambda \circ (T_t \times T_t)\, dt$$

and apply Proposition 3.11 to $(\mathcal{T} \times \mathcal{T}, \underline{\boldsymbol{\lambda}})$, $(T_1 \times T_1, \lambda)$ and the coordinate factors.

Suppose now that T_1 is 2-QS. Let $\underline{\boldsymbol{\lambda}} \in \mathcal{J}_2^e(\mathcal{T})$, $\lambda \neq \mu \otimes \mu$. There are two possibilities:

(i) $\underline{\boldsymbol{\lambda}} \in \mathcal{J}_2^e(\mathcal{T})$,

(ii) $\underline{\boldsymbol{\lambda}} \notin \mathcal{J}^e(\mathcal{T})$.

In case (i) apply Proposition 3.11 to $(\mathcal{T} \times \mathcal{T}, \underline{\boldsymbol{\lambda}})$, $(T_1 \times T_1, \lambda)$ and the coordinate factors. In case (ii) take $\lambda \in \mathcal{J}_2^e(T_1)$ such that $\underline{\boldsymbol{\lambda}} = \int_0^1 \lambda \circ (T_t \times T_t)\, dt$ and apply Proposition 3.11 to $(\mathcal{T} \times \mathcal{T}, \underline{\boldsymbol{\lambda}})$, $(T_1 \times T_1, \lambda)$ and the coordinate factors. This ends the proof. □

Proposition 3.18. *Let $l \geq 1$ and let T be an automorphism of $(X, \mathcal{B}, \mu)$, such that T^l is ergodic. Then T is 2-QS if and only if T^l is 2-QS.*

Proof. The proof goes along the same lines as the proof of Proposition 3.17. It suffices to replace integrals by finite averages and apply Proposition 3.12 instead of Proposition 3.11. □

3.4.2 Quasi-simplicity – special case

We will give now an independent proof of the fact that 2-fold simplicity of an ergodic time automorphism of a flow implies that the whole flow is 2-fold quasi-simple.

Lemma 3.19. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be an ergodic flow on $(X, \mathcal{B}, \mu)$. Consider*

$$S_t \colon (X, \mathcal{B}, \mu) \otimes ([0,1), \lambda_{[0,1)}) \to (X, \mathcal{B}, \mu) \otimes ([0,1), \lambda_{[0,1)})$$

which is a suspension over T_1. Then $\mathcal{S}$ is isomorphic to a skew product of $\mathcal{T}$ with a rotation on the circle.

Proof. Let the flow $\mathcal{S} = (S_t)_{t\in\mathbb{R}} \colon X \times [0,1) \to X \times [0,1)$ be given by

$$S_t(x, r) = (T_t y, \{r + t\}).$$

Then the map $\Psi \colon X \times [0,1) \to X \times [0,1)$ given by $\Psi \colon (x, r) \mapsto (T_r x, r)$ yields an isomorphism between $\mathcal{T}$ and $\mathcal{S}$: $\Psi \circ T_t = S_t \circ \Psi$. □

Proposition 3.20. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be an measure-preserving flow on $(X, \mathcal{B}, \mu)$ with T_1 ergodic. If T_1 is 2-fold simple then $\mathcal{T}$ is 2-QS. In fact, for every $\underline{\boldsymbol{\xi}} \in \mathcal{J}_2^e(\mathcal{T}) \setminus \mathcal{J}_2^e(T_1)$ the flow $(\mathcal{T} \times \mathcal{T}, \underline{\boldsymbol{\xi}})$ is isomorphic to a suspension flow over (T_1, μ).*

Proof. Consider $\underline{\boldsymbol{\xi}} \in \mathcal{J}_2^e(\mathcal{T}) \setminus \{\mu \otimes \mu\}$. There exists $\xi \in \mathcal{J}_2^e(T_1)$ such that $\underline{\boldsymbol{\xi}} = \int_0^1 \xi \circ (T_t \times T_t)\, dt$. Without loss of generality we may assume that this is the ergodic decomposition of $\underline{\boldsymbol{\xi}}$ for $T_1 \times T_1$. Then $\underline{\boldsymbol{\xi}}$-almost every point of the space $X \times X$ is of the form $(T_r x, T_r S x)$, where $r \in [0,1)$ and $S \in C(T_1)$. Define $\Phi \colon X \times X \to X \times \mathbb{T}$ for such points by

$$\Phi(T_r x, T_r S x) := (x, r).$$

This map is 1-1 $\underline{\boldsymbol{\xi}}$-almost everywhere. Moreover, for Borel sets $A \subset X$, $I \subset \mathbb{T}$ we have

$$\mu \otimes \lambda_{\mathbb{T}}(A \times I) = \mu(A) \cdot \lambda_{\mathbb{T}}(I).$$

On the other hand

$$\Phi^{-1}(A \times I) = \{(T_r x, T_r S x) \colon x \in A, r \in I\},$$

whence by the definition of $\underline{\boldsymbol{\xi}}$ we obtain $\underline{\boldsymbol{\xi}}(\Phi^{-1}(A \times I)) = \mu(A) \cdot \lambda_{\mathbb{T}}(I)$. Let $S_t \colon X \times \mathbb{T} \to X \times \mathbb{T}$ be given by $S_t(x, r) = (T_{[t+r]} x, \{r + t\})$. Then

$$\Phi \circ (T_t \times T_t) = S_t \circ \Phi \text{ for all } t \in \mathbb{R},$$

i.e. $\mathcal{T} \times \mathcal{T} \simeq \mathcal{S}$. The claim follows by Lemma 3.19. □

3.4.3 Distal simplicity

Proposition 3.21. *Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be a measure-preserving flow on $(X, \mathcal{B}, \mu)$ with T_1 ergodic. Then $\mathcal{T}$ is 2-DS if and only if T_1 is 2-DS.*

Proof. In the proof of Proposition 3.17 replace "Proposition 3.11" with "Proposition 3.13". □

Proposition 3.22. *Let $l \geq 1$ and let T be an automorphism of $(X, \mathcal{B}, \mu)$, such that T^l is ergodic. Then T is 2-DS if and only if T^l is 2-DS.*

Proof. In the proof of Proposition 3.18 replace "Proposition 3.12" with "Proposition 3.14". □

4 (Non)-uniqueness of the embedding

4.1 Uniqueness

We will discuss now the problem of the uniqueness of the embedding of automorphisms into measurable flows (an the uniqueness of compositional roots of automorphisms) from the point of view of self-joining properties.

Remark 4.1. Given a flow $\mathcal{T}$ the following conditions are equivalent:

(i) for any flow $\mathcal{S}$ the condition $T_1 = S_1$ implies $\mathcal{T} = \mathcal{S}$,

(ii) for any flow $\mathcal{S}$ such that $T_1 \simeq S_1$, the isomorphism between $T_1 \simeq S_1$ is an isomorphism between $\mathcal{T}$ and $\mathcal{S}$.

Remark 4.2. For weakly mixing flows $\mathcal{T}$, $\mathcal{S}$ such that $T_1 \not\perp S_1$, we always have $\mathcal{T} \not\perp \mathcal{S}$. Indeed, every non-trivial joining between T_1 and S_1 lifts to a non-trivial joining between $\mathcal{T}$ and $\mathcal{S}$. This holds in particular when $T_1 = S_1$.

Proposition 4.1. *Let $\mathcal{T}$ be a weakly mixing, measure-preserving flow on $(X, \mathcal{B}, \mu)$ with the MSJ property. Then T_1 has a unique embedding.*

We will give now two independent proofs and later on a third one.

Proof 1. Let $\mathcal{S} = (S_t)_{t\in\mathbb{R}}$ on $(X, \mathcal{B}, \mu)$ be such that $T_1 = S_1$. Then $T_1 = S_1$ is 2-fold simple by Proposition 3.1. It follows from Proposition 3.20 that $\mathcal{S}$ is 2-QS. Since by Remark 4.2 $\mathcal{T} \not\perp \mathcal{S}$, they have a non-trivial common factor (see [18]). Since $\mathcal{T}$ has the MSJ property this factor is equal to $\mathcal{B}$ and there exists $\mathcal{C} \subset \mathcal{B}$ invariant under $\mathcal{S}$, such that $\mathcal{S}|_{\mathcal{C}} = \mathcal{T}$. On the other hand, $T_1 = S_1$ which implies $\mathcal{C} = \mathcal{B}$ and the claim follows. □

Before we give the second proof, let us recall a classical result on group homomorphisms.

Theorem 4.2 (Mackey, see [80])**.** *Let G be a locally compact, second countable group and let J be a second countable group. Let $\pi\colon G \to J$ be a measurable group homomorphism. Then π is continuous.*

Proof 2 of Proposition 4.1. Let $\mathcal{S}$ be such that $T_1 = S_1$. Then for any $t \in \mathbb{R}$ we have $S_t \in C(S_1) = C(T_1) = C(\mathcal{T})$ (the latter equality follows from Proposition 3.1). By the assumption that $\mathcal{T}$ has the MSJ property, it follows that there exists a function $a\colon \mathbb{R} \to \mathbb{R}$ such that $S_t = T_{a(t)}$. It is a group homomorphism. We claim that it is continuous. Indeed, $t \mapsto S_t$ is measurable, whence (by Theorem 4.2) it is continuous. Now, using the fact that $S_t = T_{a(t)}$ and the open mapping theorem we obtain that $t \mapsto a(t)$ is also continuous. Therefore we have $a(t) = at$ for some $a \in \mathbb{R}$. By taking $t = 1$ we obtain that $a = 1$, i.e. $T_t = S_t$ for all $t \in \mathbb{R}$. □

Proposition 4.3. *If T is a weakly mixing automorphism with the MSJ property then for $k \in \mathbb{N}$ the automorphism T^k has no other roots of order k than T.*

Proof. Both above proofs of Proposition 4.1 can be easily adjusted to the case of automorphisms. □

We will show now a more universal way of proving the uniqueness of embedding. It will be used in the third proof of Proposition 4.1 and in its analogue for flows with the R-property.

Proposition 4.4. *Let $\mathcal{T}$ be a weakly mixing flow on $(X, \mathcal{B}, \mu)$ such that for any weakly mixing flow $\mathcal{S}$ on $(Y, \mathcal{C}, \nu)$ and any $\lambda \in \mathcal{J}^e(\mathcal{T}, \mathcal{S})$ either $\lambda = \mu \otimes \nu$ or the fibers in the extension $(\mathcal{T} \times \mathcal{S}, \lambda) \to (\mathcal{S}, \nu)$ are finite. Then for any $\lambda \in \mathcal{J}^e(\mathcal{T}, \mathcal{S})$, the flow $(\mathcal{T} \times \mathcal{S}, \lambda)$ is weakly mixing.*

Proof. Let $\lambda \in \mathcal{J}^e(\mathcal{T}, \mathcal{S})$. If $\lambda = \mu \otimes \nu$ then $(\mathcal{T} \times \mathcal{S}, \lambda)$ is clearly weakly mixing. Consider the case $\lambda \neq \mu \otimes \nu$ and suppose that $(\mathcal{T} \times \mathcal{S}, \lambda)$ is not weakly mixing. Denote by $\mathcal{K}$ the Kronecker factor of $(\mathcal{T} \times \mathcal{S}, \lambda)$. Since $(\mathcal{T} \times \mathcal{S}, \lambda)$ is not weakly mixing, it has $\mathcal{B} \otimes \mathcal{K}$ as a factor.[33] Consider the following disintegrations of measures:

$$\lambda = \int \mu_x \, d\mu, \quad \lambda = \int \lambda_\omega \, d\lambda|_{\mathcal{B}\otimes\mathcal{K}}, \quad \lambda|_{\mathcal{B}\otimes\mathcal{K}} = \int \widetilde{\lambda}_z \, d\mu.$$

For $f \in L^1(X \times Y, \mathcal{B} \otimes \mathcal{C}, \lambda)$ we have

$$\int f \, d\lambda = \iint f \, d\mu_x \, d\mu.$$

On the other hand,

$$\int f \, d\lambda = \iint f \, d\lambda_\omega \, d\lambda|_{\mathcal{B}\otimes\mathcal{K}} = \iiint f \, d\lambda_\omega \, d\widetilde{\lambda}_z \, d\mu = \iiint f \, d\widetilde{\lambda}_z \, d\lambda_\omega \, d\mu.$$

Since $(\mathcal{T} \times \mathcal{S}, \lambda)$ is not weakly mixing, the discrete part of the spectrum forms a countable (infninite) subgroup of $\mathbb{R}$ and the corresponding factor acts by rotations on the dual group to this group. The measures $\widetilde{\lambda}_z$ are equal to the Haar measure on this group, in particular they have infinite supports. It follows by the uniqueness of the disintegration, that also the supports of the measures μ_x have infinite number of points. This is however impossible by our assumption λ is a finite extension of μ, i.e. the measures μ_x are discrete with a finite number of atoms. The claim follows. □

Remark 4.3. The claim of the above proposition can be strenghtened, using the same arguments: If $\mathcal{T}$ is a weakly mixing flow such that for any weakly mixing flow $\mathcal{S}$ and any $\lambda \in \mathcal{J}^e(\mathcal{T}, \mathcal{S})$ the flow $(\mathcal{T} \times \mathcal{S}, \lambda)$ is also weakly mixing then any finite group extension of $\mathcal{T}$ has the same property.

Proposition 4.5. *Let $\mathcal{T}$ be a weakly mixing flow with the following property: for any weakly mixing flow $\mathcal{S}$ and any $\lambda \in \mathcal{J}^e(\mathcal{T}, \mathcal{S})$ the flow $(\mathcal{T} \times \mathcal{S}, \lambda)$ is weakly mixing. Then T_1 has a unique embedding.*

Proof. Suppose that there exists $\mathcal{S}$ such that $T_1 = S_1$ and consider the diagonal joining of $\Delta \in \mathcal{J}^e(T_1, S_1)$. We claim that $\Delta \in \mathcal{J}(\mathcal{T}, \mathcal{S})$. Indeed, otherwise for some $k \in \mathbb{N}$

$$\underline{\boldsymbol{\Delta}} := \int_0^{1/k} \Delta \circ (T_t \times S_t) \, dt \in \mathcal{J}^e(\mathcal{T}, \mathcal{S}),$$

[33] Remark that the symbol $\otimes$ used here does not mean that $\mathcal{K} \subset \mathcal{C}$. By $\mathcal{B} \otimes \mathcal{K}$ we denote the sub-σ-algebra of $\mathcal{B} \otimes \mathcal{C}$ generated by $\mathcal{B}$ and by the Kronecker factor $\mathcal{K}$. The action of $\mathcal{T} \times \mathcal{S}$ restricted to $\mathcal{B} \otimes \mathcal{K}$ is isomorphic to the Cartesian product of $\mathcal{T}$ and $(\mathcal{T} \times \mathcal{S})|_{\mathcal{K}}$.

where the measures $\Delta \circ (T_t \times S_t)$ are pairwise orthogonal for $t \in [0, 1/k)$. Then by Remark 2.3 on page 11, $(\mathcal{T} \times \mathcal{S}, \underline{\mathbf{\Delta}})$ is not weakly mixing. This yields a contradition with our assumption and ends the proof. □

Corollary 4.6. *Let $\mathcal{T}$ be a weakly mixing flow with the R-property. Then T_1 has a unique embedding.*

Proof. To see that the claim holds it suffices to combine Proposition 4.4, Proposition 4.5 and Proposition 2.6 on page 15. □

Proposition 4.7. *Let $\mathcal{T}$ be a weakly mixing flow with the MSJ property. Then for any weakly mixing flow $\mathcal{S}$ and any $\lambda \in \mathcal{J}^e(\mathcal{T}, \mathcal{S})$ other than the product measure the extension $(\mathcal{T} \times \mathcal{S}, \lambda) \to (\mathcal{S}, \nu)$ has finite fibers.*

Before we begin the proof, let us recall a result on the absence of disjointness and state a necessary lemma.

Theorem 4.8 ([46])**.** *Let $\mathcal{T}$ and $\mathcal{S}$ be ergodic flows. If $\mathcal{T}$ and $\mathcal{S}$ are not disjoint then there exists $\lambda \in \mathcal{J}^e_\infty(\mathcal{T})$ such that $(\mathcal{T}^{\times\infty}, \lambda)$ and $\mathcal{S}$ have a common factor.*

Lemma 4.9. *Let $\mathcal{T}, \widetilde{\mathcal{T}}, \mathcal{R}$ be ergodic flows. Suppose that the extension $\widetilde{\mathcal{T}} \to \mathcal{T}$ is isometric ("k:1"). Let $\widetilde{\rho} \in \mathcal{J}^e(\widetilde{\mathcal{T}}, \mathcal{R})$ and let ρ be the restriction of $\widetilde{\rho}$ to a joining between $\mathcal{T}$ and $\mathcal{R}$. Then also the extension $(\widetilde{\mathcal{T}} \times \mathcal{R}, \widetilde{\rho}) \to (\mathcal{T} \times \mathcal{R}, \rho)$ is isometric ("k:1").*

Proof. Notice that is suffices to show that the claim remains true if we replace the words "isometric extension" with the words "compact group extension". This is however clear, as for any compact group extension $\mathcal{T}_\varphi$, we can write $\mathcal{T}_\varphi \vee \mathcal{R}$ as $(\mathcal{T} \vee \mathcal{R})_{\widetilde{\varphi}}$, where $\widetilde{\varphi}(x, y) = \varphi(x)$. In this way we obtain the first part of the claim (about isometric extensions). The more detailed part (about the extensions being "k:1") follows from the fact that the group G where φ and $\widetilde{\varphi}$ take their values is the same. Moreover, any isometric extension with finite fibers is an intermediate extension for some finite group extension. □

Proof of Proposition 4.7. Let $\lambda \in \mathcal{J}^e(\mathcal{T}, \mathcal{S})$ be different from the product measure. It follows from the results in [19] that there exists $n \in \mathbb{N}$ such that the following holds:

- $\mathcal{T}^{\odot n}$ is a factor of $\mathcal{S}$,[34]
- the restriction of λ to a joining $\mathcal{T} \vee \mathcal{T}^{\odot n}$ between $\mathcal{T}$ and $\mathcal{T}^{\odot n}$ is such that $\mathcal{T} \vee \mathcal{T}^{\odot n}$ is a factor of $\mathcal{T}^{\times n}$.

Since the extension $\mathcal{T}^{\times n} \to \mathcal{T}^{\odot n}$ has finite fibers, it follows that also the intermediate extension $\mathcal{T} \vee \mathcal{T}^{\odot n} \to \mathcal{T}^{\odot n}$ has finite fibers (see the proof of Proposition 4.4). Now, applying Lemma 4.9, we obtain that the fibers in the extension $\mathcal{T} \vee \mathcal{S} \to \mathcal{S}$ are finite as well (the role of $\widetilde{\mathcal{T}}$ from Lemma 4.9 is played by $\mathcal{T} \vee \mathcal{T}^{\odot n}$, the role of $\mathcal{R}$ is played by $\mathcal{S}$ and the role of ρ is played by the joining associated with the factoring map). □

Proof 3 of Proposition 4.1. It suffices to combine Proposition 4.4, Proposition 4.5 and Proposition 4.7. □

[34] By $\mathcal{T}^{\odot n}$ we denote the symmetric factor of $\mathcal{T}^{\times n}$, i.e. the sub-σ-algebra of sets invariant under all permutations of coordinates. For the details we refer the reader e.g. to [19].

4.2 Non-uniqueness

Since the R-property implies the 2-QS property, one could ask if a result analogous to Corollary 4.6 holds also in a more general setting of 2-QS flows. It turns out this is not true. In fact, we "loose" the uniqueness of the embedding of the time-one map already in case of 2-fold simple flows.

Let D_6 stand for the smallest non-abelian group, i.e. dihedral group of order 6. Recall that the group operations in D_6 can be summarized using Cayley table as follows:

$*$	**e**	**a**	**b**	**c**	**d**	**f**
e	e	a	b	c	d	f
a	a	e	d	f	b	c
b	b	f	e	d	c	a
c	c	d	f	e	a	b
d	d	c	a	b	f	e
f	f	b	c	a	e	d

(32)

Proposition 4.10. *There exists an automorphism which is 2-fold simple and has two non-isomorphic square roots (one of these roots is 2-fold simple, the other one is only 2-QS).*

Proof. Let T be a weakly mixing automorphism with the MSJ property and consider $\varphi\colon X \to D_6$ such that $T_\varphi\colon X \times D_6 \to X \times D_6$ is also weakly mixing. Then T_φ is 2-fold simple. For $g \in D_6$ let $\sigma_g\colon X \times D_6 \to X \times D_6$ be given by $\sigma_g(x,h) = (x,hg)$. Notice that for all $g \in D_6$ we have $\sigma_g \in C(T_\varphi)$. From now on we will be using the notation from (32) for elements of D_6. Since $a^2 = e$, we have $(\sigma_a)^2 = Id$, whence

$$(T_\varphi \circ \sigma_a)^2 = (T_\varphi)^2. \tag{33}$$

Moreover, since $a * b \neq b * a$, $\sigma_b \notin C(\sigma_a)$, whence $\sigma_b \notin C(T_\varphi \circ \sigma_a)$. Therefore

$$C(T_\varphi \circ \sigma_a) \neq C(T_\varphi). \tag{34}$$

By Proposition 3.1, we have

$$C(T_\varphi) = C((T_\varphi)^2). \tag{35}$$

combining (33), (34) and (35) we conclude that

$$C((T_\varphi \circ \sigma_a)^2) \neq C(T_\varphi \circ \sigma_a).$$

It follows by Lemma 3.1 that T_φ is 2-fold simple, whereas $T_\varphi \circ \sigma_a$ is not 2-fold simple (it is 2-QS by Proposition 3.18). This ends the proof. □

Proposition 4.11. *There exists a 2-fold simple automorphism which is embeddable into two non-isomorphic flows (one of these flows is 2-fold simple and the other one is only 2-QS).*

Proof. We will follow a similar path as in the proof of Proposition 4.10. Let $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ be a weakly mixing flow on $(X, \mathcal{B}, \mu)$ with the MSJ property. Let $\varphi\colon X \times \mathbb{R} \to SU(2)$ be an $\mathbb{R}$-cocycle such that $\mathcal{T}_\varphi = ((T_t)_\varphi)_{t\in\mathbb{R}}$ is weakly mixing. Then $\mathcal{T}_\varphi$ is 2-fold simple. For $g \in SU(2)$ let $\sigma_g(x,h) = (x,hg)$ and let $\mathcal{R} = (R_t)_{t\in\mathbb{R}}$ be given by $R_t = \sigma_{g_t}$, where

$$g_t = \begin{pmatrix} e^{2\pi i t} & 0 \\ 0 & e^{-2\pi i t} \end{pmatrix}.$$

Notice that the time-one maps of $\mathcal{T}_\varphi$ and $\mathcal{T}_\varphi \circ \mathcal{R}$ are the same.[35] Moreover, for all $g \in SU(2)$ we have $\sigma_g \in C(\mathcal{T}_\varphi)$. In particular, $\sigma_{h_0} \in C(\mathcal{T}_\varphi)$, where

$$h_0 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

On the other hand, $\sigma_{h_0} \notin C(\mathcal{R})$. Indeed, we have

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} e^{2\pi it} & 0 \\ 0 & e^{-2\pi it} \end{pmatrix} = \begin{pmatrix} 0 & -e^{-2\pi it} \\ e^{2\pi it} & 0 \end{pmatrix}, \tag{36}$$

whereas

$$\begin{pmatrix} e^{2\pi it} & 0 \\ 0 & e^{-2\pi it} \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & -e^{2\pi it} \\ e^{-2\pi it} & 0 \end{pmatrix}. \tag{37}$$

Arguing as in the proof of Proposition 4.10, we conclude that

$$C(\mathcal{T}_\varphi \circ \mathcal{R}) \neq C(\mathcal{T}_\varphi) = C((T_1)_\varphi) = C((T_1 R_1)_\varphi).$$

It follows that $\mathcal{T}$ is 2-fold simple whereas $\mathcal{T} \circ \mathcal{R}$ is not 2-fold simple (it is 2-QS by Proposition 3.17). □

Remark 4.4. Notice that $(T_{1/2})_\varphi$ described in the proof of Proposition 4.11 has the properties demanded in Proposition 4.10. Indeed, substitute $t = 1/4$ into equalities (36) and (37) and use the same arguments as in the proof of Proposition 4.11.

Remark 4.5. It follows from the example presented in the proof of Proposition 4.11 that the answer to the following question is negative:

> Suppose that the extension $\widetilde{\mathcal{T}} \to \mathcal{T}$ is such that
> $\widetilde{T}_1 \to T_1$ is a compact group extension. Is is true that also
> $\widetilde{\mathcal{T}} \to \mathcal{T}$ is a compact group extension?

Similarly, it follows from the example from the proof of Proposition 4.10 that also the answer to the following question is negative:

> Suppose that the extension $\widetilde{T} \to T$ is such that
> $\widetilde{T}^2 \to T^2$ is a compact group extension. Is is true that also
> $\widetilde{T} \to T$ is a compact group extension?

Indeed, if in the constructed examples , all the considered extensions were compact group extensions, then $T_\varphi \circ \sigma_a$ and $\mathcal{T} \circ \mathcal{R}$ would be 2-fold simple, which is not the case.

Remark 4.6. In view of Remark 4.3, it is not by coincidence that for the construction from Proposition 4.11 we used a compact group extension by an infinite group.

[35] Given two commuting flows $\mathcal{T}$, $\mathcal{S}$, we denote by $\mathcal{T} \circ \mathcal{R}$ the $\mathbb{R}$-action $(T_t \circ S_t)_{t \in \mathbb{R}}$.

5 Self-joining properties of $\mathcal{T}$ and T_1 – part II

A direct consequence of Proposition 4.10 and Proposition 4.11 is that Proposition 3.20 cannot be reversed. We will now discuss this problem in more details, providing more counter-examples. Let us however see first, that the counter-examples we were looking for cannot be obtained by taking factors of products of simple systems. In particular, products of horocycle flows, their factors and their weakly mixing distal extensions do not have the desired properties.

Proposition 5.1. *Let $\mathcal{T}$ be a 2-fold simple flow with T_1 ergodic. Then any factor $\mathcal{A}$ has the following property:*

- *either $\mathcal{T}|_{\mathcal{A}}$ and $T_1|_{\mathcal{A}}$ are both 2-fold simple*
- *or $\mathcal{T}|_{\mathcal{A}}$ and $T_1|_{\mathcal{A}}$ are both 2-QS, but not 2-fold simple.*

Proof. Notice that $\mathcal{T}$ and T_1 have the same factors. Moreover, every factor corresponds to a compact subgroup of $C(\mathcal{T}) = C(T_1)$. If this subgroup is normal, then $\mathcal{T}$ and T_1 restricted to the corresponding factor are 2-fold simple. Otherwise they are only 2-QS but not 2-fold simple. □

Corollary 5.2. *Let $\mathcal{T}^{(i)}$ be 2-fold simple, weakly mixing flows. And let $\mathcal{A}$ be a factor of $\mathcal{T}^{(1)} \times \mathcal{T}^{(2)} \times \dots$ which is 2-QS. Then*

- *either $(\mathcal{T}^{(1)} \times \mathcal{T}^{(2)} \times \dots)|_{\mathcal{A}}$ and $(T_1^{(1)} \times T_1^{(2)} \times \dots)|_{\mathcal{A}}$ are both 2-fold simple,*
- *or $(\mathcal{T}^{(1)} \times \mathcal{T}^{(2)} \times \dots)|_{\mathcal{A}}$ and $(T_1^{(1)} \times T_1^{(2)} \times \dots)|_{\mathcal{A}}$ are both 2-QS, but not 2-fold simple.*

Proof. This follows immediately from Proposition 5.1 and from the fact that whenever a 2-fold QS system is a factor of a product of two weakly mixing systems, then it is a factor of one of the coordinate factors [18]. □

Denote by (P) the following property of $\mathcal{T}$:

$$T_1 \text{ is simple and } \mathcal{T} \text{ is only 2-QS.}$$

Corollary 5.3. *Let $\mathcal{T}$ be a weakly mixing flow which has the JP property, satisfying the following condition: for any factor $\mathcal{A}$ of $\mathcal{T}$ there exists a factor $\mathcal{C} \subset \mathcal{A}$ such that $\mathcal{T}|_{\mathcal{C}}$ has the (P) property. Then $\mathcal{T}$ is disjoint from weakly mixing extensions of simple flows.*

Proof. Recall [45] that a JP system is disjoint from a weakly mixing simple system if and only if they do not have a common non-trivial factor. Now it suffices to apply Corollary 5.2 and a result from [26] on lifting disjointntess. □

Remark 5.1. The assumptions of the above corollary are satisfied when $\mathcal{T}$ satisfies the (P) property and has no non-trivial factors.

5.1 Easy counter-example

Del Junco and Rudolph [19] recall an example by Glasner [29] of a weakly mixing group extension of a 2-fold simple $\mathbb{Z}$-action, which is itself not 2-fold simple. It is of the form $R(x, z_1, z_2) = (Tx, \varphi(x)z_1, z_1z_2)$, where T is an ergodic automorphism with minimal self-joinings and $\varphi\colon X \to \mathbb{T}$ yields a weakly mixing group extension T_φ of T. Del Junco and Rudolph proved that this automorphism is not 2-fold simple, by exhibiting its ergodic self-joining which is "2:1". We will follow a similar scheme and give an example of a $\mathbb{Z}_2$-extension of a 2-fold simple map such that the obtained automorphism is "2:1" and its square is 2-fold simple. We will identify $\mathbb{Z}_2$ with $\{0, 1\}$ with addition modulo 2.

Let us first recall some necessary lemmas and tools. Since we will use them for abelian groups only, we state them in such setting.

Lemma 5.4 ([19])**.** *Let T_φ be an ergodic compact group extension of a 2-fold simple automorphism T. Then T_φ is 2-fold simple if and only if for every $S \in C(T)$ there exists $\widetilde{S} \in C(T_\varphi)$ which is an extension of S.*

Lemma 5.5 ($\mathbb{Z}$-actions [54], abelian groups [49], general case [70])**.** *Suppose that T is 2-fold simple and let T_φ be its compact group extension, i.e. $T_\varphi\colon X \times G \to X \times G$, $T_\varphi(x, g) = (Tx, \varphi(x) + g)$. Let $\widetilde{S} \in C(T_\varphi)$. Then there are a continuous group automorphism $v\colon G \to G$, a measurable map $f\colon X \to G$ and $S \in C(T)$ such that $\widetilde{S}(x, g) = (Sx, f(x) + v(g))$. Equivalently, $\varphi(x) - v(\varphi(x)) = f(Tx) - f(x)$.*

Remark 5.2. If $G = \mathbb{Z}_2$ then $v(i) = i$ and $f = const$, i.e. any $\widetilde{S} \in C(T_\varphi)$ is of the form $\widetilde{S}(x, g) = (Sx, g)$ or $\widetilde{S}(x, g) = (Sx, g + 1)$ for some $S \in C(T)$.

Lemma 5.6 (see e.g. [36])**.** *Let T_φ be an ergodic compact group extension of T. Then $c \in \mathbb{T}$ is an eigenvalue of T_φ if and only if there exists a measurable function $f\colon X \to \mathbb{T}$ and a character $\chi \in \hat{G}$ such that*

$$\chi \circ \varphi = c \cdot \frac{f \circ T}{f}.$$

Let $T\colon (X, \mathcal{B}, \mu) \to (X, \mathcal{B}, \mu)$ be weakly mixing and have the MSJ property, and let $\varphi\colon X \to \mathbb{Z}_2 = \{0, 1\}$ yield a weakly mixing group extension T_φ. Consider $\psi\colon X \times \mathbb{Z}_2 \to \mathbb{Z}_2$ given by the formula $\psi(x, s) = s$. Let

$$\overline{T}(x, s, r) := (T_\varphi)_\psi(x, s, r) = (Tx, \varphi(x) + s, s + r).$$

Lemma 5.7. *$\overline{T}$ is weakly mixing.*

Proof. Suppose that $\overline{T}$ is not weakly mixing. By Lemma 5.6 there exists $c \in \mathbb{T}\setminus\{1\}$, a measurable function $\xi\colon X \times \mathbb{Z}_2 \to \mathbb{T}$ and $\chi \in \widehat{\mathbb{Z}}_2$ satisfying the equation

$$\chi \circ \psi = c \cdot \xi \circ T_\varphi / \xi. \tag{38}$$

There are two possibilities:

(a) $\chi \equiv 1$,

(b) $\chi(s) = (-1)^s$.

In case (a) equation (38) is of the form $\xi \circ T_\varphi = \overline{c} \cdot \xi$, i.e. $\overline{c}$ is an eigenvalue of T_φ. This is contradictory as T_φ is weakly mixing. In case (b) equation (38) is of the form $(-1)^s = c \cdot \xi \circ T_\varphi / \xi$. Hence $\overline{c}^2 \cdot \xi^2 = \xi^2 \circ T_\varphi$ which implies $c^2 = 1$ and $\xi^2 = 1$ almost everywhere as T_φ is weakly mixing (i.e. we have $|\xi| = 1$ almost everywhere). Now, there are again two possibilities:

(b1) $c = 1$,

(b2) $c = -1$.

Equation (38) yields

$$\begin{aligned}(-1)^s &= \xi \circ T_\varphi / \xi \text{ in case (b1)},\\ (-1)^{s+1} &= \xi \circ T_\varphi / \xi \text{ in case (b2)},\end{aligned}$$

i.e. we have

$$\begin{aligned}(-1)^s \xi(x,s) &= \xi \circ T_\varphi(x,s) = \xi(Tx, \varphi(x) + s) \text{ in case (b1)},\\ (-1)^{s+1} \xi(x,s) &= \xi \circ T_\varphi(x,s) = \xi(Tx, \varphi(x) + s) \text{ in case (b2)}.\end{aligned}$$

Using the theory of characters (ξ is a square-integrable function)

$$\xi(x,s) = \xi_1(x) + \xi_2(x)(-1)^s$$

for some $\xi_j \colon X \to \mathbb{C}$, $j = 1, 2$. Therefore

$$\begin{aligned}(-1)^s(\xi_1(x) + \xi_2(x) \cdot (-1)^s) &= \xi_1(Tx) + \xi_2(Tx) \cdot (-1)^{\varphi(x)+s} \text{ in case (b1)},\\ (-1)^{s+1}(\xi_1(x) + \xi_2(x) \cdot (-1)^s) &= \xi_1(Tx) + \xi_2(Tx) \cdot (-1)^{\varphi(x)+s} \text{ in case (b2)},\end{aligned}$$

whence

$$\begin{aligned}\xi_2(x) - \xi_1(Tx) &= (-1)^s(-\xi_1(x) + \xi_2(Tx) \cdot (-1)^{\varphi(x)}) \text{ in case (b1)},\\ -\xi_2(x) - \xi_1(Tx) &= (-1)^s(\xi_1(x) + \xi_2(Tx) \cdot (-1)^{\varphi(x)}) \text{ in case (b2)}.\end{aligned}$$

Therefore (the above equations hold for all s)

$$\begin{aligned}\xi_2(x) &= \xi_1(Tx) \text{ a for almost every } x \text{ in case (b1)},\\ \xi_2(x) &= -\xi_1(Tx) \text{ a for almost every } x \text{ in case (b2)}.\end{aligned}$$

This implies

$$\begin{aligned}\xi_1(x) &= \xi_2(Tx) \cdot (-1)^{\varphi(x)} = \xi_1(T^2x) \cdot (-1)^{\varphi(x)} \text{ in case (b1)},\\ \xi_1(x) &= -\xi_2(Tx) \cdot (-1)^{\varphi(x)} = \xi_1(T^2x) \cdot (-1)^{\varphi(x)} \text{ in case (b2)},\end{aligned}$$

whence $|\xi_1|$ is constant. In either case we have

$$\frac{\xi_1(x) \cdot \xi_1(Tx)}{\xi_1(Tx) \cdot \xi_1(T^2x)} = \frac{\xi_1(x)}{\xi_1(T^2x)} = (-1)^{\varphi(x)},$$

which means that $\chi \circ \varphi$ is a coboundary. By Lemma 5.6, this yields a contradiction as T_φ is ergodic. □

Proposition 5.8. *$\overline{T}$ is not 2-fold simple, whereas $\overline{T}^2$ is 2-fold simple.*

Proof. We will show first that $\overline{T}$ is not 2-fold simple. By Lemma 5.4 it suffices to find an element of $C(T_\varphi)$ which cannot be lifted to an element of $C(\overline{T})$. We claim that $\sigma(x,s) = (x, s+1)$ is such an automorphism. Suppose that we can find $\overline{\sigma} \in C(\overline{T})$ which projects down to σ. By Lemma 5.5, we can solve the following equation:

$$F \circ T_\varphi - F = \psi \circ \sigma - v \circ \psi,$$

where $v \in Aut(\mathbb{Z}_2)$ and $F\colon X \times \mathbb{Z}_2 \to \mathbb{Z}_2$ is measurable. Since the operations are in $\mathbb{Z}_2$, we obtain

$$\begin{aligned} F \circ T_\varphi(x,s) - F(x,s) &= \psi \circ \sigma(x,s) + \psi(x,s) \\ &= \psi(x,s+1) + \psi(x,s) = s+1+s = 1. \end{aligned}$$

We apply to the left-hand side and the right-hand side the character $\chi(r) = (-1)^r$:

$$\frac{(-1)^F \circ T_\varphi}{(-1)^F} = -1,$$

which is not possible as T_φ is weakly mixing.

We will now show that $\overline{T}^2$ is 2-fold simple. We have $\overline{T}^2 = ((T_\varphi)^2)_{\psi+\psi\circ T_\varphi}$. Since T_φ is 2-fold simple, Proposition 3.1 implies that $(T_\varphi)^2$ is also 2-fold simple and by Lemma 5.4 it suffices to show that all elements of $C((T_\varphi)^2)$ can be lifted to $C(\overline{T}^2)$. By Proposition 3.1 on page 17 we have $C((T_\varphi)^2) = C(T_\varphi)$. Therefore (see Remark 5.2) it suffices to show that $\sigma(x,s) = (x, s+1)$ lifts to an element of $C(\overline{T}^2)$, i.e. we need to find a measurable solution $F\colon X \times \mathbb{Z}_2 \to \mathbb{Z}_2$ for the following cocycle equation:

$$\psi^{(2)} \circ \sigma(x,s) + \psi^{(2)}(x,s) = F \circ (T_\varphi)^2(x,s) + F(x,s), \tag{39}$$

where $\psi^{(2)} = \psi + \psi \circ T_\varphi$. Equation (39) yields

$$\begin{aligned} F \circ (T_\varphi)^2(x,s) + F(x,s) &= \psi^{(2)}(x,s+1) + \psi^{(2)}(x,s) \\ &= \psi(x,s+1) + \psi(T_\varphi(x,s+1)) + \psi(x,s) + \psi(T_\varphi(x,s)) \\ &= s+1+\varphi(x)+s+1+s+\varphi(x)+s = 0. \end{aligned}$$

It suffices to take $F \equiv 0$, which completes the proof. □

5.2 Advanced counter-example

As already mentioned, all the counter-examples we gave until now have a major "flaw": they are not disjoint from 2-fold simple systems. We will construct a 2-QS flow which is disjoint from simple flows and whose time-one map is 2-fold simple.

5.2.1 (C, F)-constructions

Following [12] (with some minor changes corresponding to the fact that our setting is more general than the one appearing until now in the literature), we recall here the (C, F)-construction of *funny rank-one* actions.

Let G be a unimodular l.c.s.c. group. Given two subsets $E, F \subset G$, by EF we mean their product, i.e., $EF = \{ef \colon e \in E, f \in F\}$. The set $\{e^{-1} \colon e \in E\}$ is denoted by E^{-1}. If E is a singleton, say $E = \{e\}$, we write eF for EF. To define a (C, F)-action of G we need two sequences $(F_n)_{n\geq 0}$ and $(C_n)_{n>0}$ satisfying additional conditions:

$$(F_n)_{n\geq 0} \text{ is a Følner sequence in } G, \tag{40}$$
$$C_n \text{ is finite and } \#C_n > 1, \tag{41}$$
$$F_n C_{n+1} \subset F_{n+1}, \tag{42}$$
$$F_n c \cap F_n c' = \emptyset \text{ for all } c \neq c' \in C_{n+1}. \tag{43}$$

We put $X_n := F_n \times \prod_{k>n} C_k$, endow X_n with the standard Borel product σ-algebra and define a Borel embedding $X_n \to X_{n+1}$ by setting

$$(f_n, c_{n+1}, c_{n+2}, \dots) \mapsto (f_n c_{n+1}, c_{n+2}, \dots). \tag{44}$$

It is well-defined due to (42). Then we have $X_1 \subset X_2 \subset \dots$. Hence $X := \cup_n X_n$ endowed with the natural Borel σ-algebra, say $\mathcal{B}$, is a standard Borel space. Given a Borel subset $A \subset F_n$, we put

$$[A]_n := \{x \in X \colon x = (f_n, c_{n+1}, c_{n+2}, \dots) \in X_n \text{ and } f_n \in A\}$$

and call this set a *n-cylinder*. It is clear that the σ-algebra $\mathcal{B}$ is generated by the family of all cylinders.

Now we are going to define a "canonical" measure on $(X, \mathcal{B})$. Let κ_n stand for the equidistribution on C_n and $\nu_n := (\#C_1 \dots \#C_n)^{-1} \lambda_G|_{F_n}$ on F_n. We define a product measure μ_n on X_n by setting $\mu_n = \nu_n \times \kappa_{n+1} \times \kappa_{n+2} \times \dots$, $n \in \mathbb{N}$. Then the embeddings (44) are measure preserving. Hence a σ-finite measure μ on X is well defined by the restrictions $\mu|_{X_n} = \mu_n$, $n \in \mathbb{N}$. Since

$$\mu_{n+1}(X_{n+1}) = \frac{\nu_{n+1}(F_{n+1})}{\nu_{n+1}(F_n C_{n+1})} \mu_n(X_n) = \frac{\lambda_G(F_{n+1})}{\lambda_G(F_n)\#C_{n+1}} \mu_n(X_n),$$

it follows that μ is finite if and only if

$$\prod_{n=0}^{\infty} \frac{\lambda_G(F_{n+1})}{\lambda_G(F_n)\#C_{n+1}} < \infty, \text{ i.e. } \sum_{n=0}^{\infty} \frac{\lambda_G(F_{n+1} \setminus (F_n C_{n+1}))}{\lambda_G(F_n)\#C_{n+1}} < \infty. \tag{45}$$

From now on we will assume that (45) is satisfied. Moreover, we normalize λ_G in such a way that $\mu(X) = 1$.

To construct a μ-preserving action of G on (X, μ), we fix a filtration $K_1 \subset K_2 \subset \dots$ of G by compact sets. Thus $\cup_{m\geq 1} K_m = G$. Additionally assume that

$$\text{for all } m \geq 1 \text{ we have } K_m K_m \subset K_{m+1}. \tag{46}$$

Given $n, m \in \mathbb{N}$ set

$$D_m^{(n)} := \left(\bigcap_{k \in K_m} (k^{-1} F_n) \cap F_n \right) \times \prod_{l>n} C_l \subset X_n.$$

It is easy to verity that $D_{m+1}^{(n)} \subset D_m^{(n)} \subset D_m^{(n+1)}$. We define a Borel mapping

$$K_m \times D_{m+1}^{(n)} \ni (g, x) \mapsto T_{m,g}^{(n)} x \in D_m^{(n)}$$

by setting for $x = (f_n, c_{n+1}, c_{n+2}, \dots)$

$$T^{(n)}_{m,g}(f_n, c_{n+1}, c_{n+2}, \dots) := (gf_n, c_{n+1}, c_{n+2}, \dots).$$

Indeed, for $g \in K_m$ and $f_n \in \bigcap_{k\in K_{m+1}}(k^{-1}F_n) \cap F_n$ we obtain that $gf_n \in F_n$ and

$$gf_n \in \bigcap_{k\in K_{m+1}} gk^{-1}F_n = \bigcap_{k\in K_{m+1}g^{-1}} k^{-1}F_n \subset \bigcap_{k\in K_m} k^{-1}F_n$$

since by (46) we have $K_{m+1}g^{-1} \supset K_m$. Now let $D_m := \bigcup_{n=1}^{\infty} D^{(n)}_m$. Then a Borel mapping

$$T_{m,g}\colon K_m \times D_{m+1} \ni (g, x) \mapsto T_{m,g}x \in D_m$$

is well defined by the restrictions $T_{m,g}|_{D^{(n)}_m} = T^{(n)}_{m,g}$ for $g \in K_m$ and $n \geq 1$. It is clear that $D_m \supset D_{m+1}$ and $T_{m,g}|_{D_{m+2}} = T_{m+1,g}$ for all m. If follows from (40) that $\mu_n(D^{(n)}_m) \to 1$ as $n \to \infty$. Hence $\mu(D_m) = 1$ for all $m \in \mathbb{N}$. Finally we set $\widehat{X} := \bigcap_{m\geq 1} D_m$ and define a Borel mapping $T\colon G \times \widehat{X} \ni (g, x) \mapsto T_g x \in \widehat{X}$ by setting $T_g x := T_{m,g}s$ for some (and hence any) m such that $g \in K_m$. It is clear that $\mu(\widehat{X}) = 1$. Thus, we obtain that $\mathfrak{T} = (T_g)_{g\in G}$ is a free Borel measure preserving action of G on a conull subset of the standard probability space $(X, \mathcal{B}, \mu)$. It is easy to see that $\mathfrak{T}$ does not depend on the choice of filtration $(K_m)_{m\geq 1}$. We will not distinguish between two measurable sets (or mappings) which agree almost everywhere.

Definition 5.1. $\mathfrak{T}$ is called the (C, F)-action of G associated with $(C_n, F_n)_n$.

We now list some basic properties of $(X, \mu, \mathfrak{T})$. Given Borel subsets $A, B \subset F_n$ we have

$$[A \cap B]_n = [A]_n \cap [B]_n,\ [A \cup B]_n = [A]_n \cup [B]_n,$$

$$[A]_n = [AC_{n+1}]_{n+1} = \bigsqcup_{c\in C_{n+1}} [Ac]_{n+1},$$

$$T_g[A]_n = [gA]_n \text{ if } gA \subset F_n, \tag{47}$$

$$\mu([A]_n) = \#C_{n+1} \cdot \mu([Ac]_{n+1}) \text{ for every } c \in C_{n+1}, \tag{48}$$

$$\mu([A]_n) = \frac{\lambda_G(A)}{\lambda_G(F_n)}\mu(X_n), \tag{49}$$

where $\sqcup$ stands for the union of mutually disjoint sets.

We recall now the definition of funny rank one (see e.g. [22] and [23]).

Definition 5.2. Let $\mathfrak{T}$ be a measure preserving action of G on a σ-finite measure space $(X, \mathcal{B}, \mu)$.

(i) A *Rokhlin tower* or *column* for $\mathfrak{T}$ is a triple (Y, f, F), where $Y \in \mathcal{B}$, F is a relatively compact subset of G and $f\colon Y \to F$ is a measurable mapping such that for any Borel subset $H \subset F$ and an element $g \in G$ with $gH \subset F$, one has $f^{-1}(gH) = T_g f^{-1}(H)$.

(ii) We say that $\mathfrak{T}$ is of *funny rank-one* if there exists a sequence of Rokhlin towers (Y_n, f_n, F_n) such that $\lim_{n\to\infty} \mu(Y_n) = 1$ and for any subset $B \in \mathcal{B}$, there is a sequence of Borel subsets $H_n \subset F_n$ such that

$$\lim_{n\to\infty} \mu(B \triangle f_n^{-1}(H_n)) = 0.$$

It is easy to see that any funny rank-one action is ergodic. Each (C, F)-action is of funny rank-one. Note that what we call (after Danilenko) funny rank-one is called rank-one by del Junco and Yassawi in case G is discrete and countable and $G \neq \mathbb{Z}$ [21].

Proposition 5.9. *$\mathfrak{T} = (T_g)_{g\in G}$ is of funny rank-one.*

Proof. According to the definition of a funny rank-one system, we have to find a sequence of Rokhlin towers "approximating" our dynamical system. Let s_n denote the projection of $X_n = F_n \times C_{n+1} \times \dots$ onto the first coordinate. Then (X_n, s_n, F_n) satisfies the needed condition. □

5.2.2 Construction of a counter-example

In this section we will provide an example of a weakly mixing flow $\mathcal{T} = (T_t)_{t\in\mathbb{R}}$ with T_1 which is 2-fold simple such that $\mathcal{T}$ is only 2-QS (i.e. not 2-fold simple). We will do it by constructing a special (C, F)-action. Moreover, we will show later that our flow has a factor with the same self-joining properties, which is additionally disjoint from all 2-fold simple flows.

Let SU(2) stand for the special unitary group of order 2, i.e. the group of matrices with complex entries which are of the form $\begin{pmatrix} z & -\overline{\omega} \\ \omega & \overline{z} \end{pmatrix}$ and whose determinant is equal to one. Let $\lambda_{\mathrm{SU}(2)}$ denote the normalized Haar measure on SU(2). Let

$$\varphi\colon \mathbb{R} \to \mathrm{Aut}(\mathrm{SU}(2), \lambda_{\mathrm{SU}(2)})$$

be given by

$$\varphi_t(N) = \begin{pmatrix} e^{\pi i t/2} & 0 \\ 0 & e^{-\pi i t/2} \end{pmatrix} N \begin{pmatrix} e^{-\pi i t/2} & 0 \\ 0 & e^{\pi i t/2} \end{pmatrix}.$$

We consider a semidirect product $G = \mathbb{R} \ltimes_\varphi \mathrm{SU}(2)$. Recall that the multiplication in G is given by

$$(t, M)(s, N) = (t + s, M\varphi_t(N)).$$

Notice that

$$C(G) = \{(2n, I), (2n, -I)\colon n \in \mathbb{Z}\}.$$

Given $a > 0$ we let

$$\begin{aligned} I^{\mathbb{Z}}[a] &:= \{m \in \mathbb{Z}\colon |m| < a\}, \\ I^{\mathbb{Z}}_{+}[a] &:= I^{\mathbb{Z}}[a] \cup \{a\}, \\ I^{\mathbb{R}}[a] &:= (-a, a]. \end{aligned}$$

Let $(r_n)_{n\geq 0}$ be an increasing sequence of positive integers such that

$$\lim_{n\to\infty} \frac{n^4}{r_n} = 0. \tag{50}$$

We will impose later one more restriction on the growth of $(r_n)_{n\geq 0}$. We define recurrently two other sequences $(a_n)_{n\geq 0}$ and $(\widetilde{a}_n)_{n\geq 0}$ by settting

$$\begin{aligned} &a_0 = \widetilde{a}_0 = 1, \\ &a_{n+1} := \widetilde{a}_n(2r_n - 1), \\ &\widetilde{a}_{n+1} := a_{n+1} + (2n+1)\widetilde{a}_n \end{aligned} \tag{51}$$

For each $n \in \mathbb{N}$ we let

$$\begin{aligned} &H := \mathbb{Z},\ H_n := I^{\mathbb{Z}}[r_n], \\ &F_n := I^{\mathbb{R}}[a_n] \times \mathrm{SU}(2),\ \widetilde{F}_n := I^{\mathbb{R}}[\widetilde{a}_n] \times \mathrm{SU}(2), \\ &S_n := I^{\mathbb{R}}[(2n-1)\widetilde{a}_{n-1}] \times \mathrm{SU}(2). \end{aligned} \tag{52}$$

We also consider a homomorphism $\phi_n \colon H \to \mathbb{Z} \ltimes \mathrm{SU}(2) \subset G$ given by

$$\phi_n(h) = (2h\widetilde{a}_n, I).$$

Then we have

$$S_n \subset F_n,\ F_n S_n = S_n F_n \subset \widetilde{F}_n \subset G, \tag{53}$$

$$F_{n+1} = \bigsqcup_{h \in H_n} \widetilde{F}_n \phi_n(h) = \bigsqcup_{h \in H_n} \phi_n(h) \widetilde{F}_n, \tag{54}$$

$$S_{n+1} = \bigsqcup_{h \in I^{\mathbb{Z}}[n+1]} \widetilde{F}_n \phi_n(h) = \bigsqcup_{h \in I^{\mathbb{Z}}[n+1]} \phi_n(h) \widetilde{F}_n. \tag{55}$$

Fix a sequence $\varepsilon_n \to 0$ as $n \to \infty$.

For any two finite sets A, B and a map $\phi \colon A \to B$ we define a probability measure on B:

$$\mathrm{dist}_{a \in A} \phi(a) := \frac{1}{\#A} \sum_{a \in A} \delta_{\phi(a)}.$$

Given two measures κ, ρ on a finite set B let

$$\|\kappa - \rho\|_1 := \sum_{b \in B} |\kappa(b) - \rho(b)|.$$

For $n \in \mathbb{N}$ let $\widehat{S}_n$ be as in Proposition 6.6 and such that its assertion holds for sets A', B' which are sums of translations of at most $\#C_1 \cdot \ldots \cdot \#C_{n-1}$ elements form $\mathcal{A}$ (see the remark after Proposition 6.6).

Lemma 5.10 ([11], see also [16]). *If r_n is sufficiently large then there exists a map $s_n \colon H_n \to \widehat{S}_n$ such that for any $\delta \geq n^{-2} r_n$,*

$$\left\| \mathrm{dist}_{t \in I^{\mathbb{Z}}[\delta]} (s_n(h+t), s_n(h'+t), s_n(h''+t)) - \lambda_{\widehat{S}_n} \otimes \lambda_{\widehat{S}_n} \otimes \lambda_{\widehat{S}_n} \right\|_1 < \varepsilon_n \tag{56}$$

whenever $h, h', h'' \in H_n$ are pairwise distinct and $\{h, h', h''\} + I^{\mathbb{Z}}[\delta] \subset H_n$.

From now on we will assume that r_n is large enough, so that the conclusion of Lemma 5.10 is satisfied. For every $n \in \mathbb{N}$ we fix a map s_n whose existence is asserted in the lemma. Without loss of generality we may assume that the following boundary condition holds:

$$s_n(r_n - 1) = s_n(-r_n + 1) = 0.$$

Now we can define a map $c_{n+1} \colon H_n \to F_{n+1}$ by setting $c_{n+1}(h) := s_n(h)\phi_n(h)$. We also put $C_{n+1} := c_{n+1}(H_n)$.

It is easy to derive from (53) and (54) that (40), (41), (42) and (43) are satisfied for the sequence $(F_n, C_{n+1})_{n \geq 0}$. Moreover,

$$\frac{\lambda(F_{n+1})}{\lambda(F_n)\#C_{n+1}} = \frac{2a_{n+1}}{2a_n(2r_n - 1)} = \frac{\widetilde{a}_n}{a_n} = 1 + \frac{(2n-1)\widetilde{a}_{n-1}}{a_n} = 1 + \frac{2n-1}{2r_{n-1} - 1}. \tag{57}$$

From this and from (50) we deduce that (45) holds. Hence we can consider the associated (C, F)-action $\mathfrak{T}$ of G on a standard probability space $(X, \mathcal{B}, \mu)$.

Auxiliary lemmas

Lemma 5.11. *Let $f = f'\phi_{n-1}(h)$, $\hat{f} = \hat{f}'\phi_{n-1}(\hat{h})$ with $f', \hat{f}' \in \widetilde{F}_{n-1}$ and $h, \hat{h} \in H$.*

(i) Then we have

$$\begin{aligned}\widetilde{F}_{n-1}\phi_{n-1}(h + I^{\mathbb{Z}}[n-1]) &\subset fS_n \cap \hat{f}S_n \\ &\subset fS_n \cup \hat{f}S_n \subset \widetilde{F}_{n-1}\phi_{n-1}(h + I^{\mathbb{Z}}[n+1]),\end{aligned}$$

whence

$$\begin{aligned}\lambda(fS_n \setminus \widetilde{F}_{n-1}\phi_{n-1}(h + I^{\mathbb{Z}}[n-1])) &\le 4\lambda(\widetilde{F}_{n-1}), \\ \lambda(\hat{f}S_n \setminus \widetilde{F}_{n-1}\phi_{n-1}(h + I^{\mathbb{Z}}[n-1])) &\le 4\lambda(\widetilde{F}_{n-1}),\end{aligned}$$

and

$$\lambda(fS_n \triangle \hat{f}S_n) \le 4\lambda(\widetilde{F}_{n-1}).$$

(ii) If, in addition, $fS_n \subset F_n$ then

$$\frac{\lambda(AC_n \cap fS_n)}{\lambda(S_n)} = \lambda_{F_{n-1}}(A) + \overline{o}_n(1).$$

for any subset $A \subset F_{n-1}$.

Proof. By (55) we have

$$fS_n = f'\phi_{n-1}(h)\widetilde{F}_{n-1}\phi_{n-1}(I^{\mathbb{Z}}[n]) = f'\widetilde{F}_{n-1}\phi_{n-1}(h + I^{\mathbb{Z}}[n]).$$

Since $\widetilde{F}_{n-1}\widetilde{F}_{n-1} \subset \bigsqcup_{|i|\le 1} \widetilde{F}_{n-1}\phi_{n-1}(i)$, there exists a partition of $\widetilde{F}_{n-1}$ into subsets A_i, $|i| \le 1$ such that $f'A_i \subset \widetilde{F}_{n-1}\phi_{n-1}(i)$ for any i. Therefore

$$\begin{aligned}fS_n &= \bigsqcup_{|i\le 1|} f'A_i\phi_{n-1}(-i)\phi_{n-1}(i + h + I^{\mathbb{Z}}[n]) \\ &\subset \bigsqcup_{|i|\le 1} f'A_i\phi_{n-1}(-i)\phi_{n-1}(h + I^{\mathbb{Z}}[n+1]) = \widetilde{F}_{n-1}\phi_{n-1}(h + I^{\mathbb{Z}}[n+1]) \quad (58)\end{aligned}$$

since $\bigsqcup_{|i|\le 1} f'A_i\phi_{n-1}(-i) = \widetilde{F}_{n-1}$. In a similar way we obtain

$$fS_n \supset \widetilde{F}_{n-1}\varphi_{n-1}(h + I^{\mathbb{Z}}[n-1]). \quad (59)$$

Clearly, (58) and (59) remain true if we replace f with $\widehat{f}$. The remaining assertions of (i) are direct consequences of (58) and (59).

We will now prove (ii). If $fS_n \subset F_n$ then $K := h + I^{\mathbb{Z}}[n-1] \subset H_{n-1}$ and,

using (i) we obtain

$$\begin{aligned}
\frac{\lambda(AC_n \cap fS_n)}{\lambda(S_n)} &= \frac{\lambda(AC_n \cap \widetilde{F}_{n-1}\phi_{n-1}(h + I^{\mathbb{Z}}[n-1])) \pm 4\lambda(\widetilde{F}_{n-1})}{\lambda(S_n)} \\
&= \frac{1}{\lambda(S_n)} \Big(\lambda(AC_n \cap \widetilde{F}_{n-1}\phi_{n-1}(K)) \pm 4\lambda(\widetilde{F}_{n-1})\Big) \\
&= \frac{1}{\lambda(S_n)} \sum_{h' \in H_{n-1}} \Big(\lambda(As_{n-1}(h')\phi_{n-1}(h') \cap \widetilde{F}_{n-1}\phi_{n-1}(K)) \pm 4\lambda(\widetilde{F}_{n-1})\Big) \\
&= \frac{1}{\lambda(S_n)} \sum_{k \in K} \Big(\lambda(As_{n-1}(h')) \pm 4\lambda(\widetilde{F}_{n-1})\Big) \\
&= \frac{\#K}{\lambda(S_n)}\lambda(A) \pm \frac{\lambda(\widetilde{F}_{n-1})}{\lambda(S_n)} \\
&= \lambda_{F_{n-1}}(A) \cdot \frac{\lambda(F_{n-1})}{\lambda(\widetilde{F}_{n-1})} \cdot \frac{\lambda(\widetilde{F}_{n-1})}{\lambda(S_n)} \cdot \#K \pm \frac{\lambda(\widetilde{F}_{n-1})}{\lambda(S_n)} \\
&= \lambda_{F_{n-1}}(A) \cdot \frac{a_{n-1}}{\widetilde{a}_{n-1}} \cdot \frac{\widetilde{a}_{n-1} \cdot (2n-3)}{(2n-1) \cdot \widetilde{a}_{n-1}} \pm 4\frac{\widetilde{a}_{n-1}}{(2n-1)\widetilde{a}_{n-1}} \\
&= \lambda_{F_{n-1}}(A) + \overline{o}_n(1).
\end{aligned}$$

□

Lemma 5.12. *Let (G, λ) be a locally compact unimodular group with Haar measure λ. For any measurable sets $A, B, F, S \subset G$ of finite measure λ we have*

$$\int_{S\times S} \lambda(Av \cap Bw \cap F)\ dv\ dw = \int_{A\times B} \lambda(aS \cap bS \cap F)\ da\ db.$$

Proof. We will show first that for any measurable set $C \subset G$ of finite measure we have

$$\int_S \lambda(Cv \cap F)\ dv = \int_C \lambda(xS \cap F)\ dx.$$

Indeed,

$$\begin{aligned}
\int_S \lambda(Cv \cap F)\ dv &= \int_S \lambda(C \cap Fv^{-1})\ dv = \int_S \int_C \mathbb{1}_{Fv^{-1}}(x)\ dx\ dv \\
&= \int_S \int_C \mathbb{1}_{x^{-1}F}(v)\ dx\ dv = \int_C \int_S \mathbb{1}_{x^{-1}F}(v)\ dv\ dx \\
&= \int_C \lambda(S \cap x^{-1}F)\ dx = \int_C \lambda(xS \cap F)\ dx.
\end{aligned}$$

Therefore

$$\begin{aligned}
\int_{S\times S} \lambda(Av \cap Bw \cap F)\ dv\ dw &= \int_S \int_S \lambda(Av \cap (Bw \cap F))\ dv\ dw \\
&= \int_S \int_A \lambda(aS \cap (Bw \cap F))\ da\ dw = \int_A \int_S \lambda(Bw \cap (aS \cap F))\ dw\ da \\
&= \int_A \int_B \lambda(bS \cap aS \cap F)\ db\ da = \int_{A\times B} \lambda(aS \cap bS \cap F)\ da\ db.
\end{aligned}$$

□

5.2.3 Weak mixing

Proposition 5.13. *The transformation $T_{(2,I)}$ (and hence $T_{(1,I)}$) is weakly mixing.*

Proof. Let $g_n = \phi_n(1) = (2\widetilde{a}_n, I) = (2, I)^{\widetilde{a}_n}$. We will show that $(g_n)_{n\in\mathbb{N}}$ is mixing for T, i.e.

$$\lim_{n\to\infty} \mu(T_{g_n} D \cap D') = \mu(D)\mu(D') \tag{60}$$

Since $(T_g)_{g\in G}$ is of funny rank-one, it suffices to prove (60) for cylinders $D, D' \in \mathcal{A}$. Let $A = A_{(m)}$, $B = B_{(m)} \subset F_m$ be such that $A, B \in \mathcal{A}$. We will show now that

$$\mu\left(T_{g_n}\left[A\right]_m \cap \left[B\right]_m\right) = \mu\left(\left[A\right]_m\right)\mu\left(\left[B\right]_m\right) + \overline{o}_n(1),$$

where $\overline{o}_n(1)$ means (here and below) a sequence that goes to 0 as n goes to infinity and that does not depend on the choice of A, B. For $n \geq m$ we have $[A]_m = [A_{(n)}]_n$, $[B]_m = [B_{(n)}]_n$ for $A_{(n)}, B_{(n)} \subset F_n$ such that $A_{(n+1)} = A_{(n)}C_{n+1}$, $B_{(n+1)} = B_{(n)}C_{n+1}$. For $n \geq m$ let

$$\begin{aligned}
F'_n &:= F_n \cap \bigcap_{(i_1,M_1),(i_2,M_2)\in S_n} F_n(i_1, M_1)(i_2, M_2)^{-1},\\
A'_{(n)} &:= A_{(n)} \cap F'_n\\
B'_{(n)} &:= B_{(n)} \cap F'_n,\\
H'_n &:= H_n \cap (H_n - 1).
\end{aligned}$$

Notice that

$$\mu\left(\left[A_{(n)} \setminus A'_{(n)}\right]_n\right) \leq \mu\left(\left[F_n \setminus F'_n\right]_n\right)$$

and

$$\mu\left(\left[B_{(n)} \setminus B'_{(n)}\right]_n\right) \leq \mu\left(\left[F_n \setminus F'_n\right]_n\right).$$

We claim that

$$\mu\left(\left[F_n \setminus F'_n\right]_n\right) = \overline{o}_n(1). \tag{61}$$

Indeed, notice that

$$F'_n = I^{\mathbb{R}}[a_n - 2(2n-1)\widetilde{a}_{n-1}] \times \mathrm{SU}(2),$$

whence

$$\begin{aligned}
\lambda_{F_n}(F'_n) = \frac{\lambda(F'_n)}{\lambda(F_n)} &= \frac{a_n - 2(2n-1)\widetilde{a}_{n-1}}{a_n} = 1 - 2\frac{(2n-1)\widetilde{a}_{n-1}}{a_n}\\
&= 1 - 2\frac{(2n-1)\widetilde{a}_{n-1}}{\widetilde{a}_{n-1}(2r_{n-1}-1)} = 1 - 2\frac{2n-1}{2r_{n-1}-1} \to 1
\end{aligned} \tag{62}$$

by (50), and (61) follows. Therefore

$$\begin{aligned}
&\mu\left(T_{g_n}\left[A_{(n)}\right]_n \cap \left[B_{(n)}\right]_n\right) = \mu\left(T_{g_n}\left[A'_{(n)}\right]_n \cap \left[B'_{(n)}\right]_n\right) + \overline{o}_n(1)\\
&= \sum_{h\in H_m} \mu\left(T_{g_n}\left[A'_{(n)}c_{n+1}(h)\right]_{n+1} \cap \left[B'_{(n)}\right]_n\right) + \overline{o}_n(1).
\end{aligned} \tag{63}$$

Notice that $g_n \in C(G)$ for all $n \in \mathbb{N}$. Hence

$$\begin{aligned} g_n A c_{n+1}(h) &= g_n A s_n(h)\phi_n(h) = A s_n(h) g_n \phi_n(h) \\ &= A s_n(h)\phi_n(h+1) = A s_n(h) s_n(h+1)^{-1} c_{n+1}(h+1) \end{aligned}$$

provided that $h, h+1 \in H_n$. Moreover, for all $h \in H_n$

$$\mu\left(T_{g_n}\left[A'_{(n)}c_{n+1}(h)\right]_{n+1}\right) = \frac{\mu\left(\left[A'_{(n)}\right]_n\right)}{\#C_{n+1}} \leq \frac{\mu\left([F_n]_n\right)}{\#C_{n+1}} = \frac{\mu(X_n)}{\#C_{n+1}} = \overline{o}_n(1).$$

Therefore

$$\begin{aligned} &\sum_{h\in H_n} \mu\left(T_{g_n}\left[A'_{(n)}c_{n+1}(h)\right]_{n+1} \cap \left[B'_{(n)}\right]_n\right) \\ &\quad = \sum_{h\in H'_n} \mu\left(T_{g_n}\left[A'_{(n)}c_{n+1}(h)\right]_{n+1} \cap \left[B'_{(n)}\right]_n\right) \\ &\quad = \sum_{h\in H'_n} \mu\left(\left[A'_{(n)}s_n(h)s_n(h+1)^{-1}c_{n+1}(h+1)\right]_{n+1} \cap \left[B'_{(n)}\right]_n\right) \end{aligned} \tag{64}$$

Notice that

$$A'_{(n)}s_n(h)s_n(h+1)^{-1}c_{n+1}(h+1) \subset F_n c_{n+1}(h+1) \subset F_{n+1}.$$

Moreover,

$$\left[B'_{(n)}\right]_n = \bigsqcup_{\widetilde{h}\in H_n} \left[B'_{(n)}c_{n+1}(\widetilde{h})\right]_{n+1},$$

where $B'_{(n)}c_{n+1}(\widetilde{h}) \subset F_n c_{n+1}(\widetilde{h})$ and

$$\left[A'_{(n)}s_n(h)s_n(h+1)^{-1}c_{n+1}(h+1)\right]_{n+1} \cap \left[B'_{(n)}c_{n+1}(\widetilde{h})\right]_{n+1} = \emptyset$$

when $h+1 \neq \widetilde{h}$. Hence

$$\begin{aligned} &\sum_{h\in H'_n} \mu\left(\left[A'_{(n)}s_n(h)s_n(h+1)^{-1}c_{n+1}(h+1)\right]_{n+1} \cap [B'_{(n)}]_n\right) \\ &\quad = \sum_{h\in H'_n} \left(\left[\left(A'_{(n)}s_n(h)s_n(h+1)^{-1} \cap B'_{(n)}\right)c_{n+1}(h+1)\right]_{n+1}\right) \\ &\quad = \frac{1}{\#H'_n}\sum_{h\in H'_n} \mu\left(\left[A'_{(n)}s_n(h)s_n(h+1)^{-1} \cap B'_{(n)}\right]_n\right) \\ &\quad = \frac{1}{\#H'_n}\sum_{h\in H'_n} \lambda_{F_n}\left(A'_{(n)}s_n(h)s_n(h+1)^{-1} \cap B'_{(n)}\right)\mu(X_n) \\ &\quad = \frac{1}{\#H'_n}\sum_{h\in H'_n} \lambda_{F_n}\left(A'_{(n)}s_n(h) \cap B'_{(n)}s_n(h+1)\right) + \overline{o}_n(1) \\ &\quad = \frac{1}{\#H'_n}\sum_{h\in H'_n} \lambda_{F_n}\left(A_{(n)}s_n(h) \cap B_{(n)}s_n(h+1)\right) + \overline{o}_n(1). \end{aligned} \tag{65}$$

Recall that $A_{(n)} = A_{(n-1)}C_n$, $B_{(n)} = B_{(n-1)}C_n$ and $\left[A_{(n-1)}\right]_{n-1}$, $\left[B_{(n-1)}\right]_{n-1}$ are sums of at most $\#C_1 \cdot \ldots \cdot \#C_{n-1}$ translations of some elements from $\mathcal{A}$ (recall that $[A_{(m)}]_m$, $[B_{(m)}]_m \in \mathcal{A}$). Let $\xi_n := \operatorname{dist}_{h\in H'_n}(s_n(h), s_n(h+1))$. It follows from Lemma 5.10 that

$$\left\|\xi_n - \lambda_{\widehat{S}_n} \otimes \lambda_{\widehat{S}_n}\right\| < \varepsilon_n = \overline{o}_n(1).$$

Define $f\colon S_n \times S_n \to \mathbb{R}$ by

$$f(v,w) = \lambda_{F_n}\left(A_{(n)}v \cap B_{(n)}w\right).$$

Then

$$\begin{aligned}
&\frac{1}{\#H'_n}\sum_{h\in H'_n} \lambda_{F_n}\left(A_{(n)}s_n(h) \cap B_{(n)}s_n(h+1)\right)\\
&\quad= \int_{S_n\times S_n} \lambda_{F_n}\left(A_{(n)}v \cap B_{(n)}w\right)\ d\xi_n(v,w)\\
&\quad= \int f\ d\xi_n = \int f\ d\lambda_{\widehat{S}_n}\otimes\lambda_{\widehat{S}_n} \pm \varepsilon_n\\
&\quad= \int_{S_n\times S_n} \lambda_{F_n}(A_{(n-1)}C_nv \cap B_{(n-1)}C_nw)\ d\lambda_{\widehat{S}_n}\otimes\lambda_{\widehat{S}_n}(v,w) \pm \varepsilon_n.
\end{aligned} \tag{66}$$

Hence by the choice of $\widehat{S}_n$ we obtain

$$\begin{aligned}
&\int_{S_n\times S_n}\lambda_{F_n}\left(A_{(n-1)}C_nv \cap B_{(n-1)}C_nw\right)\ d\lambda_{\widehat{S}_n}\otimes\lambda_{\widehat{S}_n}(v,w)\\
&\quad= \int_{S_n\times S_n} \lambda_{F_n}\left(A_{(n-1)}C_nv \cap B_{(n-1)}C_nw\right)\ d\lambda_{S_n}\otimes\lambda_{S_n}(v,w) \pm \varepsilon_n\\
&\quad= \sum_{h,h'\in H_{n-1}} \int_{S_n\times S_n} \frac{\lambda(A_{(n-1)}c_n(h)v \cap B_{(n-1)}c_n(h')w \cap F_n)}{\lambda(F_n)}\ d\lambda_{S_n}(v)\ d\lambda_{S_n}(w) \pm \varepsilon_n\\
&\quad= \sum_{h,h'\in H_{n-1}} \int_{A_{(n-1)}\times B_{(n-1)}} \frac{\lambda(ac_n(h)S_n \cap bc_n(h')S_n \cap F_n)}{(\lambda(S_n))^2\lambda(F_n)}\ d\lambda(a)\ d\lambda(b) \pm \varepsilon_n.
\end{aligned} \tag{67}$$

For $a \in A_{(n-1)}$, $b \in B_{(n-1)}$ we have

$$ac_n(h) = as_{n-1}(h)\phi_{n-1}(h) \in \widetilde{F}_{n-1}\phi_{n-1}(h), \tag{68}$$

$$bc_n(h') = bs_{n-1}(h')\phi_{n-1}(h') \in \widetilde{F}_{n-1}\phi_{n-1}(h'). \tag{69}$$

Moreover,

$$\begin{aligned}
(0,I)c_n(h) = c_n(h) = s_{n-1}(h)\phi_{n-1}(h) \in \widetilde{F}_{n-1}\phi_{n-1}(h),\\
(0,I)c_n(h') = c_n(h') = s_{n-1}(h')\phi_{n-1}(h') \in \widetilde{F}_{n-1}\phi_{n-1}(h').
\end{aligned}$$

Therefore by Lemma 5.11 (i) we obtain

$$\begin{aligned}
&\lambda\left((ac_n(h)S_n \cap bc_n(h')S_n \cap F_n)\triangle(c_n(h)S_n \cap c_n(h')S_n \cap F_n)\right)\\
&\qquad\le \lambda(ac_n(h)S_n\triangle c_n(hS_n) + \lambda(bc_n(h')S_n\triangle c_n(h')S_n) \le 8\lambda(\widetilde{F}_{n-1}).
\end{aligned} \tag{70}$$

By (68), (69) and, again, Lemma 5.11 (i)

$$ac_n(h)S_n \cap bc_n(h')S_n \subset \widetilde{F}_{n-1}\phi_{n-1}(h + I^{\mathbb{Z}}[n+1]) \cap \widetilde{F}_{n-1}\phi_{n-1}(h' + I^{\mathbb{Z}}[n+1]).$$

Hence $ac_n(h)S_n \cap bc_n(h')S_n \neq \emptyset$ only if $h' - h \in I^{\mathbb{Z}}[2n+1]$. If the latter is satisfied, we say that h and h' are *partners*. Denote by $P(h)$ the set of partners of h from H_{n-1}. Clearly, $\#P(h) \leq 4n+1$. Therefore we deduce from (63), (64), (65), (66), (67) that

$$\begin{aligned}
&\mu(T_{g_n}[A]_m \cap [B]_m) \\
&\quad= \sum_{h\in H_{n-1}} \sum_{h'\in P(h)} \frac{\lambda(A_{(n-1)})\lambda(B_{(n-1)})}{(\lambda(S_n))^2\lambda(F_n)} \\
&\qquad\cdot \Big(\lambda(c_n(h)S_n \cap c_n(h')S_n \cap F_n) \pm 8\lambda(\widetilde{F}_{n-1})\Big) \\
&\quad= \frac{\lambda(A_{(n-1)})\lambda(B_{(n-1)})}{(\lambda(S_n))^2\lambda(F_n)} \\
&\qquad\cdot \sum_{h\in H_{n-1}} \sum_{h'\in P(h)} \lambda(c_n(h)S_n \cap c_n(h')S_n \cap F_n) \\
&\qquad\pm \lambda(H_{n-1})(4n+1)\frac{(\lambda(F_{n-1}))^2}{(\lambda(S_n))^2\lambda(F_n)} \cdot 4\lambda(\widetilde{F}_{n-1}) \\
&\quad= \frac{\lambda(A_{(n-1)})\lambda(B_{(n-1)})}{(\lambda(F_{n-1}))^2}\theta_n \\
&\qquad\pm \lambda(H_{n-1})(4n+1)\frac{(\lambda(F_{n-1}))^2}{(\lambda(S_n))^2\lambda(F_n)} \cdot 4\lambda(\widetilde{F}_{n-1}) \\
&\quad= \lambda_{F_{n-1}}(A_{(n-1)})\lambda_{F_{n-1}}(B_{(n-1)})\theta_n \\
&\qquad\pm \lambda(H_{n-1})(4n+1)\frac{(\lambda(F_{n-1}))^2}{(\lambda(S_n))^2\lambda(F_n)} \cdot 4\lambda(\widetilde{F}_{n-1}),
\end{aligned} \tag{71}$$

where $\theta_n > 0$. Notice that (57) together with (50) means that $a_n/\widetilde{a}_n \to 1$ as $n \to \infty$. Therefore, by (51) and (52) we have

$$\begin{aligned}
&\lambda(H_{n-1})(4n+1)\frac{(\lambda(F_{n-1}))^2}{(\lambda(S_n))^2\lambda(F_n)} \cdot 4\lambda(\widetilde{F}_{n-1}) \\
&\qquad= \frac{(2r_{n-1}-1)(4n+1)(2a_{n-1})^2 \cdot 4 \cdot 2\widetilde{a}_{n-1}}{((2n-1)\widetilde{a}_{n-1})^2 \cdot 2a_n} \\
&\qquad= \frac{16(4n+1)}{2n-1} \cdot \left(\frac{a_{n-1}}{\widetilde{a}_{n-1}}\right)^2 \cdot \frac{\widetilde{a}_{n-1}(2r_{n-1}-1)}{a_n(2n-1)} \\
&\qquad= \frac{16(4n+1)}{2n-1} \cdot \left(\frac{a_{n-1}}{\widetilde{a}_{n-1}}\right)^2 \cdot \frac{1}{2n-1} = \overline{o}_n(1).
\end{aligned}$$

Therefore substituting $A_{(n-1)} = B_{(n-1)} = F_{n-1}$ in (71) (see Remark 6.2) and passing to the limit we obtain $\theta_n \to 1$ as $n \to \infty$. Therefore by (49)

$$\mu(T_{g_n}[A]_m \cap [B]_m) = \mu([A]_m)\mu([B]_m) + \overline{o}_n(1) \tag{72}$$

and the claim follows. □

5.2.4 Self-joining properties

For $k = (t, M) \in G$ let $k^* = (1, I)(t, M)(1, I)^{-1} = (t, \varphi_1(M))$. Notice that $(k^*)^* = k$.

Proposition 5.14. *The transformation $T_{(1,I)}$ is "2:1". Moreover,*

$$\mathcal{J}_2^e(T_{(1,I)}) = \left\{ \frac{1}{2} \left(\mu_{T_k} + \mu_{T_{k^*}} \right) : k \in G \right\}.$$

Proof. Take any joining $\nu \in J_2^e(T_{(1,I)})$. Let $I_n := I^{\mathbb{Z}}[n^{-2}a_n]$, $J_n = I^{\mathbb{Z}}[n^{-2}r_n]$ and $\Phi_n = I_n + 2\widetilde{a}_n J_n$. We first notice that $(\Phi_n)_{n=1}^{\infty}$ is a Følner sequence in $\mathbb{Z}$. Since

$$\frac{a_n}{n^2} + \frac{2\widetilde{a}_n r_n}{n^2} < \frac{\widetilde{a}_n(2r_n + 1)}{n^2} < \frac{2a_{n+1}}{(n+1)^2},$$

it follows that $\Phi_n \subset I_{n+1} + I_{n+1}$ and hence $\cup_{m=1}^{n} \Phi_m \subset I_{n+1} + I_{n+1}$. This implies

$$\#(\Phi_{n+1} - \cup_{m=1}^{n} \Phi_m) \leq 3\#\Phi_{n+1}$$

for every $n \in \mathbb{N}$, i.e. Shulman's condition [48] is satisfied for $(\Phi_n)_{n=1}^{\infty}$. By [48], the pointwise ergodic theorem holds along $(\Phi_n)_{n=1}^{\infty}$ for any ergodic transformation. Since $T_{(1,I)}$ is ergodic by Proposition 5.13, we have

$$\frac{1}{\#\Phi_n} \sum_{i \in \Phi_n} \mathbb{1}_D(T_{(i,I)}x)\mathbb{1}_{D'}(T_{(i,I)}x') \to \nu(D \times D') \tag{73}$$

as $n \to \infty$ for ν-a.a. $(x, x') \in X \times X$ and for all cylinders $D, D' \subset X$. We call such (x, x') a generic point for $(T_{(1,I)} \times T_{(1,I)}, \nu)$. Fix one of them. Then $x, x' \in X_n$ for all sufficiently large n and we have the following expansions:

$$x = (f_n, c_{n+1}(h_n), c_{n+2}(h_{n+1}), \dots),$$
$$x' = (f'_n, c_{n+1}(h'_n), c_{n+2}(h'_{n+1}), \dots)$$

with $f_n, f'_n \in F_n$ and $h_i, h'_i \in H_i$, $i > n$. We let $H_n^- := I^{\mathbb{Z}}[(1 - n^{-2})r_n] \subset H_n$. Then

$$\frac{\#H_n^-}{\#H_n} \geq 1 - n^{-2}.$$

Since the marginals of ν are both equal to μ, by the Borel-Cantelli lemma we may assume without loss of generality that $h_n, h_n^- \in H_n^-$ for all sufficiently large n. This implies, in turn, that

$$f_{n+1} = f_n c_{n+1}(h_n) \in \widetilde{F}_n \phi_n(H_n^-) \subset I_+^{\mathbb{Z}}[(2r_n(1 - n^{-2}) - 1)\widetilde{a}_n] \times SU(2)$$

and, similarly,

$$f'_{n+1} \in I_+^{\mathbb{Z}}[(2r_n(1 - n^{-2}) - 1)\widetilde{a}_n] \times SU(2).$$

Notice that given $g \in \Phi_n$ we have $(g, I) = (b, I)\phi_n(t)$ for some uniquely determined $b \in I_n$ and $t \in J_n$. Moreover, $t + h_n \in H_n$. We also claim that

$$(b, I)f_n S_n S_n = (b, I)f_n S_n S_n^{-1} \subset F_n \tag{74}$$

To verify this, it suffices to show that

$$\frac{a_n}{n^2}+\left(2r_{n-1}\left(1-\frac{1}{(n-1)^2}\right)-1\right)\widetilde{a}_{n-1}+2(2n-1)\widetilde{a}_{n-1}<a_n.$$

We will show that the following stronger inequality holds for n is large enough:

$$\frac{a_n}{n^2}+2r_{n-1}\left(1-\frac{1}{(n-1)^2}\right)\widetilde{a}_{n-1}+4n\widetilde{a}_{n-1}<a_n,$$

i.e.

$$2r_{n-1}\left(1-\frac{1}{(n-1)^2}\right)+4n<(2r_{n-1}-1)\left(1-\frac{1}{n^2}\right).$$

The latter inequality, in turn, is equivalent to

$$1+4n-\frac{1}{n^2}<2r_{n-1}\left(\frac{1}{(n-1)^2}-\frac{1}{n^2}\right),$$

which follows from (50) in a routine way. Hence

$$\begin{aligned}(g,I)f_ns_n(h_n)\phi_n(h_n)&=dc_{n+1}(t+h_n),\\(g,I)f'_ns_n(h'_n)\phi_n(h'_n)&=d'c_{n+1}(t+h'_n),\end{aligned}$$

where

$$\begin{aligned}d:=&(b,I)f_ns_n(h_n)s_n(t+h_n)^{-1},\\d':=&(b,I)f'_ns_n(h'_n)s_n(t+h'_n)^{-1}\end{aligned}$$

belong to F_n by (74).

We consider separately two cases:

(i) $h_n=h'_n$ for all $n\geq N$. Then $f_n{f'_n}^{-1}=f_N{f'_N}^{-1}=:k$ for all $n\geq N$. Then

$$(b,I)k(b,I)^{-1}=\begin{cases}k, & \text{when } b \text{ is even}\\ k^*, & \text{when } b \text{ is odd.}\end{cases}$$

(ii) $h_n\neq h'_n$ for infinitely many, say, *bad* n.

Our goal is to prove that

$$\nu=\begin{cases}\frac{1}{2}(\mu_{T_k}+\mu_{T_{k^*}}), & \text{in case (i)}\\ \mu\otimes\mu, & \text{in case (ii)}\end{cases}$$

Notice that it suffices to show the corresponding equalities of measures restricted to the classes of cylinder sets from $\mathcal{A}$. We may also impose additional assumptions on the cylinders under consideration, provided that every set can still be approximated by finite sums of such cylinders.

Take $[A_{(m)}]_m,\ [B_{(m)}]_m\in\mathcal{A}$ and for $n\geq m$ set

$$A_{(n+1)}=A_{(n)}C_{n+1},\ B_{(n+1)}=B_{(n)}C_{n+1}.$$

We have

$$\begin{aligned}
&\frac{1}{\#\Phi_n}\#\left\{g\in\Phi_n\colon\ \left(T_{(g,I)}x,T_{(g,I)}x'\right)\in[A_{(m)}]_m\times[B_{(m)}]_m\right\}\\
&\quad=\frac{1}{\#\Phi_n}\#\left\{g\in\Phi_n\colon\ \left(T_{(g,I)}x,T_{(g,I)}x'\right)\in[A_{(n)}]_n\times[B_{(n)}]_n\right\}\\
&\quad=\frac{1}{\#I_n}\sum_{b\in I_n}\frac{\#\left\{t\in J_n\colon d\in A_{(n)},d'\in B_{(n)}\right\}}{\#J_n}\\
&\quad=\frac{1}{\#I_n}\sum_{b\in I_n}\xi_n\left(A_{(n)}^{-1}(b,I)f_ns_n(h_n)\times B_{(n)}^{-1}(b,I)f_n's_n(h_n')\right),
\end{aligned}\tag{75}$$

where

$$\xi_n=\operatorname{dist}_{t\in J_n}\left(s_n(t+h_n),s_n(t+h_n')\right).$$

We begin by treating the case (i). Take a cylinder $D'\in\mathcal{A}$ of order m which is a finite sum of cylinders from $\mathcal{A}$. It is clear that $\widetilde{D}':=D'\cap[k^{-1}F_m]_m$ is also a finite sum of cylinders from $\mathcal{A}$. Moreover,

$$\begin{aligned}
\mu(D'\triangle\widetilde{D}')&=\mu([\sqcup_{j=1}^{j_0}A_j]_m\setminus[k^{-1}F_m]_m)\\
&=\frac{\lambda(\sqcup_{j=1}^{j_0}A_j\setminus k^{-1}F_m)}{\lambda(F_m)}\mu(X_m)\le\frac{\lambda(F_m\setminus k^{-1}F_m)}{\lambda(F_m)}\\
&=\mu\left(\left[F_m\setminus k^{-1}F_m\right]_m\right).
\end{aligned}$$

Given ε for m large enough $\mu\left(\left[F_m\setminus k^{-1}F_m\right]_m\right)<\varepsilon$ since $(I^{\mathbb{R}}[a_m])_{m\in\mathbb{R}}$ is a Følner sequence in $\mathbb{R}$. Therefore we may additionally assume that every cylinder $[B]_m\in\mathcal{A}$ satisfies the condition $kB\subset F_m$ if m is large enough. Since $kF_m=k^*F_m$, it follows immediately that also $k^*B\subset F_m$.

It is easy to deduce from Lemma 5.10 that $\|\xi_n-\Delta\|_1<\varepsilon_n$ for $n>N$, where Δ is the probability equidistributed on $\widehat{S}_n\times\widehat{S}_n$. This yields (having in mind how we have chosen $\widehat{S}_n$)

$$\begin{aligned}
&\frac{1}{\#I_n}\sum_{b\in I_n}\xi_n\left(A_{(n)}^{-1}(b,I)f_ns_n(h_n)\times B_{(n)}^{-1}(b,I)f_n's_n(h_n')\right)\\
&\quad=\frac{1}{\#I_n}\sum_{b\in I_n}\lambda_{\widehat{S}_n}\left(A_{(n)}^{-1}(b,I)f_ns_n(h_n)\cap B_{(n)}^{-1}(b,I)f_n's_n(h_n)\right)\pm\varepsilon_n\\
&\quad=\frac{1}{\#I_n}\sum_{b\in I_n}\lambda_{S_n}\left(A_{(n)}^{-1}(b,I)f_ns_n(h_n)\cap B_{(n)}^{-1}(b,I)f_n's_n(h_n)\right)\pm2\varepsilon_n
\end{aligned}\tag{76}$$

We obtain further

$$\begin{aligned}
&\frac{1}{\#I_n}\sum_{b\in I_n}\lambda_{S_n}\left(A_{(n)}^{-1}(b,I)f_n s_n(h_n)\cap B_{(n)}^{-1}(b,I)f'_n s_n(h_n)\right)\\
&\quad=\frac{1}{\#I_n}\sum_{b\in I_n}\frac{\lambda\left(A_{(n)}\cap(b,I)f_n {f'}_n^{-1}(b,I)^{-1}B_{(n)}\cap(b,I)f_n s_n(h_n)S_n\right)}{\lambda(S_n)}\\
&\quad=\frac{1}{\#I_n}\sum_{2b\in I_n}\frac{\lambda\left(A_{(n)}\cap kB_{(n)}\cap(2b,I)f_n s_n(h_n)S_n\right)}{\lambda(S_n)}\\
&\qquad+\frac{1}{\#I_n}\sum_{2b+1\in I_n}\frac{\lambda\left(A_{(n)}\cap k^*B_{(n)}\cap(2b,I)f_n s_n(h_n)S_n\right)}{\lambda(S_n)}.
\end{aligned}\tag{77}$$

Since $kB_{(m)}\subset F_m$, we have $kB_{(n)}\subset F_n$. Therefore

$$A_{(n-1)}C_n\cap kB_{(n-1)}C_n=\left(A_{(n-1)}\cap kB_{(n-1)}\right)C_n.$$

It follows from Lemma 5.11 (ii) and from (74) that

$$\begin{aligned}
&\frac{\lambda\left(A_{(n)}\cap kB_{(n)}\cap(2b,I)f_n s_n(h_n)S_n\right)}{\lambda(S_n)}\\
&\qquad=\frac{\lambda\left(\left(A_{(n-1)}\cap kB_{(n-1)}\right)C_n\cap(2b,I)f_n s_n(h_n)S_n\right)}{\lambda(S_n)}\\
&\qquad=\lambda_{F_{n-1}}\left(A_{(n-1)}\cap kB_{(n-1)}\right)+\overline{o}_n(1)\\
&\qquad=\frac{\mu\left(\left[A_{(n-1)}\cap kB_{(n-1)}\right]_{n-1}\right)}{\mu(X_{n-1})}+\overline{o}_n(1)\\
&\qquad=\mu\left(\left[A_{(m)}\right]_m\cap T_k\left[B_{(m)}\right]_m\right)+\overline{o}_n(1).
\end{aligned}\tag{78}$$

In a similar way we obtain

$$\begin{aligned}
&\frac{\lambda\left(A_{(n)}\cap k^*B_{(n)}\cap(2b,I)f_n s_n(h_n)S_n\right)}{\lambda(S_n)}\\
&\qquad=\mu\left(\left[A_{(m)}\right]_m\cap T_{k^*}\left[B_{(m)}\right]_m\right)+\overline{o}_n(1).
\end{aligned}\tag{79}$$

If follows from (73), (75), (76), (77), (78) and (79) that

$$\nu=\frac{1}{2}(\mu_{T_k}+\mu_{T_{k^*}}).$$

We pass now to the second case where $h_n\neq h'_n$ for infinitely many, say, *bad* n. n. It follows from Lemma 5.10 that $\left\|\xi_n-\lambda_{\widehat{S}_n}\otimes\lambda_{\widehat{S}_n}\right\|_1<\varepsilon_n$ for all such n. Therefore

$$\begin{aligned}
&\frac{1}{\#I_n}\sum_{b\in I_n}\xi_n\left(A_{(n)}^{-1}(b,I)f_n s_n(h_n)\times B_{(n)}^{-1}(b,I)f'_n s_n(h'_n)\right)\\
&\quad=\frac{1}{\#I_n}\sum_{b\in I_n}\lambda_{\widehat{S}_n}\left(A_{(n)}^{-1}(b,I)f_n s_n(h_n)\right)\lambda_{\widehat{S}_n}\left(B_{(n)}^{-1}(b,I)f'_n s_n(h'_n)\right)\pm\varepsilon_n\\
&\quad=\frac{1}{\#I_n}\sum_{b\in I_n}\lambda_{S_n}\left(A_{(n)}^{-1}(b,I)f_n s_n(h_n)\right)\lambda_{S_n}\left(B_{(n)}^{-1}(b,I)f'_n s_n(h'_n)\right)\pm 2\varepsilon_n,
\end{aligned}\tag{80}$$

where the last equality follows from our choice of $\widehat{S}_n$. Now we derive from Lemma 5.11 (ii) that

$$\lambda_{S_n}\left(A_{(n)}^{-1}(b,I)f_n s_n(h_n)\right) = \frac{\lambda\left(A_{(n)}\cap(b,I)f_n s_n(h_n)S_n\right)}{\lambda(S_n)}$$

$$= \lambda_{F_{n-1}}\left(B_{(n-1)}\right) + \overline{o}_n(1) = \frac{\mu\left(\left[A_{(n-1)}\right]_{n-1}\right)}{\mu(X_n)} = \mu\left([A_{(m)}]_m\right) + \overline{o}_n(1) \quad (81)$$

and

$$\lambda_{S_n}\left(B_{(n)}^{-1}(b,I)f'_n s_n(h'_n)\right) = \mu\left([B_{(m)}]_m\right) + \overline{o}_n(1). \quad (82)$$

It follows from (75), (80), (81) and (82) that $\nu = \mu\otimes\mu$, which ends the proof. □

Proposition 5.15. *$T_{(1,I)}$ is 3-fold PID.*

Sketch of the proof. Let $\nu \in \mathcal{J}_3^e(T_{(1,I)})$ be pairwise independent. We will show that $\nu = \mu^{\otimes 3}$. To this end, we will follow the same path as in the proof of Proposition 5.14. Fix a generic point (x,x',x'') for $(T_{(1,I)}^{\times 3},\nu)$. Then $x,x',x''\in X_n$ for all sufficiently large n. Without loss of generality we may assume that

$$x = (f_n, c_{n+1}(h_n), c_{n+1}(h_{n+1}),\dots),$$
$$x' = (f_n, c_{n+1}(h'_n), c_{n+1}(h'_{n+1}),\dots),$$
$$x'' = (f_n, c_{n+1}(h''_n), c_{n+1}(h''_{n+1}),\dots),$$

with $f_n, f'_n, f''_n \in F_n$ and $h_i, h'_i, h''_i \in H_i^-$ for $i > n$. Now take

$$[A_{(m)}]_m, [A'_{(m)}]_m, [A''_{(m)}]_m \in \mathcal{A}$$

and set $A_{(n+1)} = A_{(n)}C_{n+1}$, $A'_{(n+1)} = A'_{(n)}C_{n+1}$, $A''_{(n+1)} = A''_{(n)}C_{n+1}$ for $n \geq m$. Then as in the proof of previous proposition we have

$$\frac{1}{\#\Phi_n}\#\left\{g\in\Psi_n\colon (T_{g,I}x, T_{g,I}x', T_{g,I}x'')\in [A_{(m)}]_m\times[A'_{(m)}]_m\times[A''_{(m)}]_m\right\}$$
$$= \frac{1}{\#I_n}\sum_{b\in I_n}\xi_n(A_{(n)}^{-1}d\times {A'_{(n)}}^{-1}d\times {A''_{(n)}}^{-1}d), \quad (83)$$

where $d = (b,I)f_n s_n(h_n)$, $d' = (b,I)f'_n s_n(h'_n)$, $d = (b,I)f''_n s_n(h''_n)$ and

$$\xi_n = dist_{t\in J_n}\left(s_n(t+h_n), s_n(t+h'_n), s_n(t+h''_n)\right).$$

Next, given $k > 0$,

$$\frac{\mu\otimes\mu\left(\{(y,y')\in X_k\times X_k\colon y_i\neq y'_i \text{ for all } i>k\}\right)}{\mu(X_k)\mu(X_k)}$$
$$= \prod_{i>k}\frac{\#C_i^2-\#C_i}{\#C_i}\prod_{i>k}\left(1-\frac{1}{\#H_i}\right) > 0,$$

where $(y_i)_{i\geq k}$, $(y'_i)_{i\geq k}$ are the "coordinates" of $y,y'\in X_k = F_k\times C_{k+1}\times C_{k+2}\times\dots$ respectively. Since ν-almost every point is generic for $(T_{(1,I)}^{\times 3},\nu)$, we can select (x,x',x'') in such a way that h_i, h'_i, h''_i are pairwise distinct for all $i\geq$

n whenever n is large enough. Therefore, it follows from Lemma 5.10 that $\|\xi_i - \lambda_{\widehat{S}_i} \otimes \lambda_{\widehat{S}_i} \otimes \lambda_{\widehat{S}_i}\| < \varepsilon_i$ for all $i \geq n$. Now we derive from (83) and Lemma 5.11 (having in mind how we have chosen the sets $\widehat{S}_i$) that

$$\nu([A_{(m)}]_m \times [A'_{(m)}]_m \times [A''_{(m)}]_m) = \mu([A_{(m)}]_m)\mu([A'_{(m)}]_m)\mu([A''_{(m)}]_m) + \overline{o}_n(1).$$

This implies $\nu = \mu^{\otimes 3}$. □

Corollary 5.16. *$T_{(1,I)}$ is PID.*

Proof. In view of the above proposition, this follows immediately by [28]. □

Remarks

1. In view of Proposition 5.14 and the definition of k^* (see page 53), one can expect that $T_{(2,I)}$ is 2-fold simple. This is indeed the case. A rigorous proof goes along the same path as the proof of Proposition 5.14 (with some simplifications). Moreover, since $T_{(1,I)}$ is PID, $T_{(2,I)}$ is also PID [11], i.e. it is simple.

2. It follows from Proposition 5.14 and Proposition 3.17 that the constructed example has as its sub-action the flow $(T_{(t,I)})_{t\in\mathbb{R}}$ which is only QS (it is not simple) with a simple time-2 map $T_{(2,I)}$. However, we cannot say at this stage yet, that the flow under consideration is disjoint from simple flows (cf. Corollary 5.3) – it is clear only that it has no factors which would be simple (see the next remark).

3. Recall that since $T_{(2,I)}$ is simple, any of its factors is determined by a compact subgroup of $C(T_{(2,I)}) = G$. The largest compact subgroup of G is $\{0\} \times SU(2) \simeq SU(2)$ and it is normal in G. It determines the smallest non-trivial factor of T_1, we will denote it by $\mathcal{A}(SU(2)) = \mathcal{A}$. Notice that $\mathcal{A}$ is also a factor of $\mathcal{T}$. Take $g \in G \setminus \{0\} \times SU(2)$. Then $\frac{1}{2}(\mu_{T_g} + \mu_{T_{g^*}})|_{\mathcal{A}\otimes\mathcal{A}}$ is a 2-fold ergodic joining of $\mathcal{T}|_{\mathcal{A}}$ and it is not difficult to see that it is "2:1" over its marginals (this argument is taken from [11]). Since $\mathcal{T}|_{\mathcal{A}}$ has no non-trivial factors, in view of Corollary 5.3 we obtain an example of a flow disjoint from simple flows, which is QS and its time-2 map is simple.

4. An easy linear time change yields now a flow $\mathcal{T}$ which is QS, it is disjoint from simple flows, and such that T_1 is simple. In particular, T_1 cannot be embedded into a simple flow.

6 Appendix – uniform distribution

6.1 Necessary tools

All the information in this section is taken from [40], unless stated otherwise. We will often use it freely, without precise reference.

Definition 6.1. Let X be a compact Hausdorff space with a (regular) probability Borel measure μ. A a sequence $(x_n)_{n\in\mathbb{N}} \subset X$ is said to be *uniformly distributed* in (X, μ) if for every continuous function $f\colon X \to \mathbb{R}$ we have

$$\lim_{N\to\infty} \frac{1}{N} \sum_{n=1}^{N} f(x_n) = \int_X f \, d\mu.$$

Definition 6.2. Let $X = \mathbb{R}^s$. We say that $(x_n)_{n\in\mathbb{N}} \subset X$ is *uniformly distributed mod* 1 if $(\{x_n\})_{n\in\mathbb{N}}$ is uniformly distributed in $[0,1)^s$ with respect to the Lebesgue measure.[36]

Definition 6.3. Let $(x_n)_{n\in\mathbb{N}} \subset \mathbb{R}^s$. Discrepancy is classically defined in the following two ways: for $N \in \mathbb{N}$ we let

$$D_N^* = D_N^*(x_1,\dots,x_N) = \sup_{J^*} \left| \frac{1}{N}\sum_{n=1}^{N} \mathbb{1}_{J^*}(\{x_n\}) - \lambda_{\mathbb{R}^s}(J^*) \right|,$$
$$D_N = D_N(x_1,\dots,x_N) = \sup_{J} \left| \frac{1}{N}\sum_{n=1}^{N} \mathbb{1}_{J}(\{x_n\}) - \lambda_{\mathbb{R}^s}(J) \right|.$$

where J^* runs over all subsets of $[0,1)^s$ of the form $[0,\beta_1)\times\cdots\times[0,\beta_s)$ and J runs over subsets of $[0,1)^s$ of the form $[\alpha_1,\beta_1)\times\cdots\times[\alpha_s,\beta_s)$.

Proposition 6.1. *Let $(x_n)_{n\in\mathbb{N}} \subset \mathbb{R}^s$. The following conditions are equivalent:*

- *the sequence $(x_n)_{n\in\mathbb{N}}$ is uniformly distributed mod* 1,
- *$D_N \to 0$ as $N \to \infty$*
- *$D_N^* \to 0$ as $N \to \infty$.*

Definition 6.4. Let $\mathcal{S} = \{(x_{n,\sigma})\colon \sigma \in J\}$ be a family of sequences in X where J denotes an arbitrary index set. $\mathcal{S}$ is called a family of *equi-μ-uniformly distributed* sequences in X if for every $f \in C(X)$, we have

$$\lim_{N\to\infty} \sup_{\sigma\in J} \left| \sum_{n=1}^{N} f(x_{n,\sigma}) - \int f \, d\mu \right| = 0.$$

Proposition 6.2 ([40])**.** *Let X be a compact Hausdorff uniform space. Suppose that $\{P_\sigma\colon \sigma\in J\}$ is a family of measure-preserving transformations on X (with respect to a given probability Borel measure μ on X) that is equicontinuous at every point $x \in X$, and let $(x_n)_{n\in\mathbb{N}}$ be a given μ-uniformly distributed sequence in X. Then $\{(P_\sigma x_n)\colon \sigma\in J\}$ is a family of equi-μ-uniformly distributed sequences in X.*

Theorem 6.3 ([34])**.** *(see also [40]) Let $(x_n)_{n\in\mathbb{N}} \subset \mathbb{R}^s$ be uniformly distributed mod* 1 *and let $f\colon [0,1]^s \to \mathbb{R}$ be a continuous function. Let*

$$D_N^*(f,(x_n)_{1\le n\le N}) := \left| \frac{1}{N}\sum_{i=1}^{N} f(x_i) - \int_{[0,1]^s} f \, d\lambda_{\mathbb{R}^s} \right|.$$

Then

$$D_N^*(f,(x_n)_{1\le n\le N}) \le (1+2^{2s-1})M\left(\left[(D_N^*(f,(x_n)_{1\le n\le N}))^{-1/k}\right]^{-1}\right),$$

where M is the modulus of continuity of f.

[36] We treat $[0,1)$ as a circle, i.e. we consider $[0,1]$ and identify the points 0 and 1.

Corollary 6.4. *Let $(x_n)_{n\in\mathbb{N}} \subset \mathbb{R}^s$ be uniformly distributed mod 1 and let $\mathcal{F}$ be a family of equi-continuous functions on $[0,1]^s$. Then*

$$\sup_{f\in\mathcal{F}} D_N^*(f,(x_n)_{n\in\mathbb{N}}) \to 0 \text{ as } N\to\infty.$$

Theorem 6.5. *(von Neumann, Oxtoby, Ulam, see [62]) Let μ_1,μ_2 be non-atomic probability Borel measures on $[0,1]^s$, with full support and such that $\mu_i(\partial([0,1]^s)) = 0$, $i=1,2$. Then there exists a homeomorphism $h\colon [0,1]^s \to [0,1]^s$ such that $\mu_1 \circ h^{-1} = \mu_2$. Moreover, $h|_{\partial([0,1]^s)} = id$.*

6.2 Technical results

By λ_G, $\lambda_{SU(2)}$ we will denote the Haar measures on G and $SU(2)$ respectively, such that $\lambda_G((0,1)\times SU(2)) = 1$ and $\lambda_{SU(2)}(SU(2)) = 1$. Given a measurable set $S\subset G$ by λ_S we denote the conditional probability measure on S. Finally, given a finite set $\widehat{S}\subset G$, by $\lambda_{\widehat{S}}$ we denote the normalized counting measure, i.e. $\lambda_{\widehat{S}}(A) = \frac{\#(\widehat{S}\cap A)}{\#\widehat{S}}$. Given $n\in\mathbb{N}$, let $\varepsilon_n > 0$.

Proposition 6.6. *There exists a dense family $\mathcal{A}$ of subsets of $G = \mathbb{R}\ltimes_\varphi SU(2)$ and for any $n\in\mathbb{N}$ there exists a finite subset $\widehat{S}_n \subset S_n$ such that the following hold:*

(i) For all $A,B\in\mathcal{A}$ and all $a,b\in G$ we have

$$\lambda_{S_n}(A^{-1}a\cap B^{-1}b) = \lambda_{\widehat{S}_n}(A^{-1}a\cap B^{-1}b) \pm \varepsilon_n.$$

(ii) For all $A,B\in\mathcal{A}$ and all $a,b\in G$ we have

$$\int_{S_n\times S_n} f d\ \lambda_{S_n}\otimes\lambda_{S_n} = \int_{S_n\times S_n} f d\ \lambda_{\widehat{S}_n}\otimes\lambda_{\widehat{S}_n} \pm \varepsilon_n$$

for $f\colon G\times G\to\mathbb{R}$ of the form $f(v,w) = \lambda_{F_n}(Aav\cap Bbw)$.

Before we begin the proof, we prepare tools and provide some necessary lemmas. First we introduce the notation. Let $\mathcal{C}_d$ stand for the family of cubes in $[0,1]^d$. By $\boldsymbol{d}\colon \mathbb{R}\times\mathbb{R}\to\mathbb{R}$ we will denote the maximum metric in $\mathbb{R}^d$ and by $\boldsymbol{d}(x,C)$ we will denote the distance of the point x from the set C. Given $C\in\mathcal{C}_d$ and $\varepsilon>0$ let

$$C_\varepsilon := \{x\notin C\colon \boldsymbol{d}(x,C) < \varepsilon\}$$

and

$$f_{C,\varepsilon} := \max\left(1 - \frac{\boldsymbol{d}(x,C)}{\varepsilon}, 0\right).$$

Notice that for each $\varepsilon>0$

$$\{f_{C,\varepsilon}\colon C\in\mathcal{C}_d\} \text{ is an equi-continuous family of functions} \tag{84}$$

and

$$\lambda(C_\varepsilon) \le 2d\varepsilon \text{ for all } C\in\mathcal{C}_d \text{ and } \varepsilon>0. \tag{85}$$

Lemma 6.7. *Let (X, μ) be a compact metric space with a probability Borel measure and let $(y_n)_{n\in\mathbb{N}} \subset X$ be uniformly distributed with respect to μ. Let $\mathcal{F}$ be a family of measure-preserving equi-continuous homeomorphisms from X to $[0,1]^d$. Then*

$$\sup_{\Psi\in\mathcal{F}} D_N^*\left((\Psi(y_n))_{1\le n\le N}\right) \to 0.$$

Proof. Let $\varepsilon > 0$ and choose $0 < \delta < \varepsilon$ small enough, so that for every cube C_δ whose sides have length at most δ we have $\Psi^{-1}(C_\delta) \subset C'_\varepsilon$, where C'_ε is some cube whose sides have length at most ε. Let $K := d \cdot 2^{d-1} \cdot \frac{1}{\delta}$.

We have

$$\left|\frac{1}{N}\sum_{n=1}^{N} \mathbb{1}_C \circ \Psi(x_n) - \lambda(C)\right| \le \underbrace{\left|\frac{1}{N}\sum_{n=1}^{N} \left(\mathbb{1}_C \circ \Psi(x_n) - f_{C,\delta} \circ \Psi(x_n)\right)\right|}_{I_1}$$
$$+ \underbrace{\left|\frac{1}{N}\sum_{n=1}^{N} f_{C,\delta} \circ \Psi(x_n) - \int f_{C,\delta}\, d\lambda\right|}_{I_2} + \underbrace{\left|\int f_{C,\delta}\, d\lambda - \lambda(C)\right|}_{I_3}.$$

Notice that

$$I_1 \le \frac{1}{N}\sum_{n=1}^{N} \mathbb{1}_{C_\delta} \circ \Psi(x_n) = \frac{1}{N}\sum_{n=1}^{N} \mathbb{1}_{\Psi^{-1}(C_\delta)}(x_n).$$

By the choice of K the set C_δ is included in a union of (at most) K cubes whose sides are equal at most δ. Therefore, by the choice of δ, $\Psi^{-1}(C_\delta)$ is included in a union of (at most) K cubes whose sides are equal at most ε. It follows immediately that there exists N_0 such that for $N \ge N_0$ we have $I_1 \le \varepsilon$.

By Theorem 6.3 we have $I_2 \to 0$. Moreover, additionally taking (84) into account, we see that the speed of convergence can be estimated using only $D_N^*((x_n)_{1\le n\le N})$ (and ε).

Finally, $I_3 \le \lambda(C_\delta) < 2d\delta \le 2d\varepsilon$ and the claim follows.

□

Lemma 6.8. *Let $(y_n)_{n\in\mathbb{N}} \subset [0,1]^d$ be uniformly distributed. Let $\mathcal{F}$ be a family of equicontinuous homeomorphisms of $[0,1]^d$ preserving the Lebesgue measure. Then*

$$\sup_{\Psi_1,\Psi_2\in\mathcal{F}} \sup_{C,D,E\subset\mathcal{C}_d} \left|\frac{1}{N}\sum_{n=1}^{N} \mathbb{1}_{\Psi_1(C)\cap\Psi_2(D)\cap E}(y_n) - \lambda(\Psi_1(C)\cap\Psi_2(D)\cap E)\right| \to 0.$$

Proof. Choose $\varepsilon > 0$ and let

$$I_1 = \left|\lambda(\Psi_1(C)\cap\Psi_2(D)\cap E) - \int f_{C,\varepsilon}\circ\Psi_1^{-1}\cdot f_{D,\varepsilon}\circ\Psi_2^{-1}\cdot f_{E,\varepsilon}\, d\lambda\right|,$$

$$I_2 = \left|\frac{1}{N}\sum_{n=1}^{N} \mathbb{1}_{\Psi_1(C)\cap\Psi_2(D)\cap E}(x_n) - \frac{1}{N}\sum_{n=1}^{N} (f_{C,\varepsilon}\circ\Psi_1^{-1}\cdot f_{D,\varepsilon}\circ\Psi_2^{-1}\cdot f_{E,\varepsilon})(x_n)\right|,$$

$$I_3 = \left|\frac{1}{N}\sum_{n=1}^{N} (f_{C,\varepsilon}\circ\Psi_1^{-1}\cdot f_{D,\varepsilon}\circ\Psi_2^{-1}\cdot f_{E,\varepsilon})(x_n) - \int f_{C,\varepsilon}\circ\Psi_1^{-1}\cdot f_{D,\varepsilon}\circ\Psi_2^{-1}\cdot f_{E,\varepsilon}\, d\lambda\right|.$$

Clearly, for the proof of lemma, it suffices to show that $I_1, I_2, I_3 \to 0$ and the convergence is uniform with respect to $C, D, E \in \mathcal{C}_d$ and $\Psi_1, \Psi_2 \in \mathcal{F}$.

We have

$$\begin{aligned}
I_1 \leq & \left| \int \mathbb{1}_C \circ \Psi_1^{-1} \cdot \mathbb{1}_D \circ \Psi_2^{-1} \cdot \mathbb{1}_E - \mathbb{1}_C \circ \Psi_1^{-1} \cdot \mathbb{1}_D \circ \Psi_2^{-1} \cdot f_{E,\varepsilon} \, d\lambda \right| \\
& + \left| \int \mathbb{1}_C \circ \Psi_1^{-1} \cdot \mathbb{1}_D \circ \Psi_2^{-1} \cdot f_{E,\varepsilon} - \mathbb{1}_C \circ \Psi_1^{-1} \cdot f_{D,\varepsilon} \circ \Psi_2^{-1} \cdot f_{E,\varepsilon} \right| \\
& + \left| \int \mathbb{1}_C \circ \Psi_1^{-1} \cdot f_{D,\varepsilon} \circ \Psi_2^{-1} \cdot f_{E,\varepsilon} - f_{C,\varepsilon} \circ \Psi_1^{-1} \cdot f_{D,\varepsilon} \circ \Psi_2^{-1} \cdot f_{E,\varepsilon} \right| \\
\leq & \int |\mathbb{1}_E - f_{E,\varepsilon}| \, d\lambda + \int \left| \mathbb{1}_D \circ \Psi_2^{-1} - f_{D,\varepsilon} \circ \Psi_2^{-1} \right| \, d\lambda \\
& + \int \left| \mathbb{1}_C \circ \Psi_1^{-1} - f_{C,\varepsilon} \circ \Psi_1^{-1} \right| \, d\lambda \\
= & \int |\mathbb{1}_E - f_{E,\varepsilon}| \, d\lambda + \int |\mathbb{1}_D - f_{D,\varepsilon}| \, d\lambda + \int |\mathbb{1}_C - f_{C,\varepsilon}| \, d\lambda \\
\leq & \lambda(E_\varepsilon) + \lambda(D_\varepsilon) + \lambda(C_\varepsilon) \leq 6d\varepsilon,
\end{aligned}$$

where the last inequality follows by (85).

We will now estimate I_2:

$$\begin{aligned}
I_2 \leq & \left| \frac{1}{N} \sum_{n=1}^{N} \left[(\mathbb{1}_{\Psi_1(C)} \cdot \mathbb{1}_{\Psi_2(D)} \cdot \mathbb{1}_E)(x_n) - (\mathbb{1}_{\Psi_1(C)} \cdot \mathbb{1}_{\Psi_2(D)} \cdot f_{E,\varepsilon}) \right] (x_n) \right| \\
& + \left| \frac{1}{N} \sum_{n=1}^{N} \left[(\mathbb{1}_{\Psi_1(C)} \cdot \mathbb{1}_{\Psi_2(D)} \cdot f_{E,\varepsilon})(x_n) - (\mathbb{1}_{\Psi_1(C)} \cdot (f_{D,\varepsilon} \circ \Psi_2^{-1}) \cdot f_{E,\varepsilon})(x_n) \right] \right| \\
& + \left| \frac{1}{N} \sum_{n=1}^{N} \left[(\mathbb{1}_{\Psi_1(C)} \cdot (f_{D,\varepsilon} \circ \Psi_2^{-1}) \cdot f_{E,\varepsilon})(x_n) - ((f_{C,\varepsilon} \circ \Psi_1^{-1}) \cdot (f_{D,\varepsilon} \circ \Psi_2^{-1}) \cdot f_{E,\varepsilon})(x_n) \right] \right| \\
\leq & \frac{1}{N} \sum_{n=1}^{N} \mathbb{1}_{E_\varepsilon}(x_n) + \frac{1}{N} \sum_{n=1}^{N} \mathbb{1}_{\Psi_2(D_\varepsilon)}(x_n) + \frac{1}{N} \sum_{n=1}^{N} \mathbb{1}_{\Psi_1(C_\varepsilon)}(x_n) \\
= & \frac{1}{N} \sum_{n=1}^{N} \mathbb{1}_{E_\varepsilon}(x_n) + \frac{1}{N} \sum_{n=1}^{N} \mathbb{1}_{D_\varepsilon}(\Psi_2^{-1}(x_n)) + \frac{1}{N} \sum_{n=1}^{N} \mathbb{1}_{C_\varepsilon}(\Psi_1^{-1}(x_n))
\end{aligned}$$

Since each of the sets $E_\varepsilon, D_\varepsilon, C_\varepsilon$ is a difference of two cubes, the above expression converges to $\lambda(E_\varepsilon) + \lambda(D_\varepsilon) + \lambda(C_\varepsilon) \leq 6d\varepsilon$. Moreover, in view of Lemma 6.7 the convergence is uniform with respect to $C, D, E \in \mathcal{C}_d$ and $\Psi_1, \Psi_2 \in \mathcal{F}$.

I_3 also converges to zero and again, in view of Theorem 6.3, the convergence is uniform with respect to $C, D, E \in \mathcal{C}_d$ and $\Psi_1, \Psi_2 \in \mathcal{F}$ which completes the proof. □

Let d be a right-invariant metric on G. Let $\mathcal{U} = \{U_i \colon 1 \leq i \leq M\}$ be a finite cover of $SU(2)$ by open sets, such that $\lambda_{SU(2)}(\partial U_i) = 0$ for $1 \leq i \leq M$. Then

$$\mathcal{V} := \bigcup_{l \in \mathbb{Z}} \bigcup_{i=1}^{M} (l-1, l+2) \times U_i \tag{86}$$

is an open cover of G, such that $\lambda_G(\partial((l-1,l+2)\times U_i))=0$.

Lemma 6.9. *The cover (86) has a positive Lebesgue number.*[37]

Proof. Suppose that there is no $\delta>0$ such that for each $(t,\boldsymbol{M})\in G$ we have $B((t,\boldsymbol{M}),\delta)\subset(l-1,l+2)\times U_i$ for some $l\in\mathbb{Z}$ and $1\le i\le M$. We claim that

$$B(x,\varepsilon)\subset(l-1,l+2)\times U_i \iff B(x(-l,I),\varepsilon)\subset(-1,2)\times U_i. \tag{87}$$

Indeed, assume that $B((t,\boldsymbol{M}),\varepsilon)\subset(l-1,l+2)\times U_i$ and take $y\in B((t,\boldsymbol{M})(-l,I),\varepsilon)$. We have $d(y(l,I),(t,\boldsymbol{M}))=d(y,(t,\boldsymbol{M})(-l,I))<\varepsilon$, whence

$$y\in[(l-1,l+2)\times U_i]\,(-l,I)=(-1,2)\times U_i$$

and (87) follows. According to our assumption there exists $x_n=(t_n,\boldsymbol{M}_n)$ such that

$$\text{none of the balls } B\left(x_n,\tfrac{1}{n}\right) \text{ is a subset of some element of our cover.} \tag{88}$$

Without loss of generality, in view of (87), we may assume that

$$x_n\in(-1,2)\times U_i \text{ for some } i$$

and also that $x_n\to x$ for some $x=(t,\boldsymbol{M})\in(l-1,l+2)\times U_j$, where $l\in\mathbb{Z}$, $1\le j\le M$. Then for N large enough we have $B(x,\frac{1}{N})\subset(l-1,l+2)\times U_j$. Let $n\ge 2N$ be large enough, so that $x_n\in B(x,\frac{1}{2N})$. Then $B(x_n,\frac{1}{n})\subset B(x_n,\frac{1}{2N})\subset B\left(x,\frac{1}{N}\right)\subset(l-1,l+2)\times U_j$, which yields a contradiction with (88). This ends the proof. □

For $1\le i\le M$ let $\psi_i\colon\overline{U}_i\to[0,1]^3$ be a homeomorphism which carries $\lambda_{\overline{U}_i}$ to the Lebesgue measure $\lambda_{[0,1]^3}$ on $[0,1]^3$.[38] For $1\le i\le M$ and $l\in\mathbb{Z}$ let $\psi_{l,i}\colon[l-1,l+2]\times\overline{U}_i\to[-1,2]\times[0,1]^3$ be given by

$$\psi_{l,i}(x,\boldsymbol{M})=(x-l,\psi_i(\boldsymbol{M}))$$

and let $\psi\colon[-1,2]\times[0,1]^3\to[0,1]^4$ be given by

$$\psi(x_1,x_2,x_3,x_4)=((x_1+1)/3,x_2,x_3,x_4)\,.$$

For $a\in G$ let $g_a,g\colon G\to G$ be given by

$$g_a(b)=ba \text{ and } g(b)=b^{-1} \text{ for } b\in G.$$

Lemma 6.10. *The family of functions*

$$\left\{g_a\circ g\circ\psi_{l,i}^{-1}\circ\psi^{-1}\colon a\in G,l\in\mathbb{Z},1\le i\le M\right\}$$

is uniformly bi-equicontinuous.

Proof. It suffices to show that the following families of functions are uniformly bi-equicontinuous:

[37] Recall that if the metric space (X,d) is compact and an open cover of X is given, then there exists a number $\delta>0$ such that every subset of X having diameter less than δ is contained in some member of the cover. Any such δ is called a Lebesgue number of this cover.

[38] Such homeomorphism exists in view of Theorem 6.5.

(i) $\mathcal{F}_1 = \left\{\psi_{l,i}^{-1} \colon l \in \mathbb{Z}, 1 \leq i \leq M\right\}$,

(ii) $\mathcal{F}_2 = \left\{g|_{[l-1,l+2]\times SU(2)} \colon l \in \mathbb{Z}\right\}$,

(iii) $\mathcal{F}_3 = \{g_a \colon a \in G\}$.

Notice that $\mathcal{F}_1$ is a finite family of homeomorphisms of compact spaces and elements of $\mathcal{F}_3$ are isometries, so both $\mathcal{F}_1, \mathcal{F}_3$ are uniformly bi-equi continuous. We consider now $\mathcal{F}_2$. For $l \in \mathbb{Z}$ let $g_l = g|_{[l-1,l+2]\times SU(2)}$. For

$$(t, M), (s, N) \in [l-1, l+2] \times SU(2)$$

we have

$$d((t-l, M), (s-l, N)) = d((t, M)(-l, I), (s, N)(-l, I)) = d((t, M), (s, N)).$$

and

$$\begin{aligned} d(g_l(t, M), g_l(s, N)) &= d((-t, \varphi_{-t}(M^{-1})), (-s, \varphi_{-s}(N^{-1}))) \\ &= d((-t, \varphi_{-t}(M^{-1}))(l, I), (-s, \varphi_{-s}(N^{-1}))(l, I)) \\ &= d((l-t, \varphi_{-t}(M^{-1})), (l-s, \varphi_{-s}(N^{-1}))) \\ &= d((l-t, \varphi_{l-t}(M^{-1})), (l-s, \varphi_{l-s}(N^{-1}))) \\ &= d(g_0(t-l, M), g_0(s-l, N)). \end{aligned}$$

Moreover, $g_l^{-1} = g_{-1-l}$. This ends the proof. □

Let $\{\rho_i\}_{1\leq i\leq M}$ be a smooth partition of unity subordinate to the cover $\mathcal{U}$ and let $\{\eta_l\}_{l\in\mathbb{Z}}$ be a smooth partition of unity subordinate to the cover $\cup_{l\in\mathbb{Z}}(l-1, l+2) = \mathbb{R}$. Then clearly $\{\eta_l \otimes \rho_i\}_{1\leq i\leq M, l\in\mathbb{Z}}$ is a smooth partition of unity subordinate to cover $\mathcal{V}$.

Let $\widetilde{\mathcal{A}}$ be a family of "cubes" in $[0,1]^4$ sufficiently small, so that for $a_1, a_2, a_3 \in G$, $l_1, l_2, l_3 \in \mathbb{Z}$, $1 \leq i_1, i_2, i_3 \leq M$, $C_1, C_2, D_1, D_2, D_3 \in \widetilde{\mathcal{A}}$ whenever the sets

$$\bigcup_{j=1}^{2} g_{a_j} \circ g \circ \psi_{l_j,i_j}^{-1} \circ \psi^{-1}(C_j) \text{ and } \bigcup_{j=1}^{3} g_{a_j} \circ \psi_{l_j,i_j}^{-1} \circ \psi^{-1}(D_j) \text{ are connected}$$

then

$$\bigcup_{j=1}^{2} g_{a_j} \circ g \circ \psi_{l_j,i_j}^{-1} \circ \psi^{-1}(C_j) \subset (l_C - 1, l_C + 2) \times U_{i_C}$$

and

$$\bigcup_{j=1}^{3} g_{a_j} \circ \psi_{l_j,i_j}^{-1} \circ \psi^{-1}(D_j) \subset (l_D - 1, l_D + 2) \times U_{i_D} \tag{89}$$

for some $l_{C,D} \in \mathbb{Z}, 1 \leq i_C, i_D \leq M$. This can be done in view of Lemma 6.9 and Lemma 6.10.

Remark 6.1. Without loss of generality, we may assume that in fact an even stronger condition than (89) holds:

$$\left(\bigcup_{j=1}^{3} g_{a_j} \circ \psi_{l_j,i_j}^{-1} \circ \psi^{-1}(D_j)\right)_{\delta_0} \cup \left(\bigcup_{j=1}^{3} g_{a_j} \circ \psi_{l_j,i_j}^{-1} \circ \psi^{-1}(D_j)\right) \subset (l_D - 1, l_D + 2) \times U_{i_D}, \tag{90}$$

where $\delta_0 > 0$ is a sufficiently small number, which is independent of the choice of D_j, a_j, $1 \leq j \leq 3$. This can be done for example by adding one more step before fixing the open cover of $SU(2)$. Namely, given $\{U_i\}_{1\leq i\leq M}$, we need to find $\delta_0 > 0$ such that $\{(U_i)_{\delta_0} \cup U_i\}_{1\leq i\leq M}$ still consists of sets homeomorphic to $(0,1)^4$ and work with both covers simultaneously.

Let

$$\mathcal{A} = \left\{\psi_{l,i}^{-1} \circ \psi^{-1}(C) \colon C \in \widetilde{\mathcal{A}}, l \in \mathbb{Z}, 1 \leq i \leq M\right\}.$$

Let $(x_n)_{n\in\mathbb{N}} \subset [0,1] \times SU(2)$ be a sequence which is uniformly distributed in $[0,1]\times SU(2)$.[39] For $l \in \mathbb{Z}$ we denote by $(x_{n,l})_{n\in\mathbb{N}}$ the sequence in $[l,l+1]\times SU(2)$ given by $x_{n,l} = x_n + (l,0)$ (the addition is understood coordinate-wise).

Proof of Proposition 6.6 (i). Fix $n \in \mathbb{N}$. The set $\widehat{S}_n$ we are looking for will consists of elements of sequences $(x_{n,l})_{n\in\mathbb{N}}$, more precisely, we will have

$$\widehat{S}_n = \{x_{n,l} \colon 1 \leq n \leq n_0, l \in \mathbb{Z}\} \cap S_n,$$

where $n_0 \in \mathbb{N}$ will be some sufficiently large number (which will depend on ε_n). Take $A, B \in \mathcal{A}$ and all $a, b \in G$. Then

$$A = \psi_{l_A,i_A}^{-1} \circ \psi^{-1}(C_A),\ B = \psi_{l_B,i_B}^{-1} \circ \psi^{-1}(C_B)$$

for some $C_A, C_B \in \widetilde{\mathcal{A}}$ and $l_A, l_B \in \mathbb{Z}$, $1 \leq i_A, i_B \leq M$.

If $A^{-1}a \cap B^{-1}b \cap S_n = \emptyset$ then clearly

$$\lambda_{S_n}(A^{-1}a \cap B^{-1}b) = \lambda_{\widehat{S}_n}(A^{-1}a \cap B^{-1}b) = 0,$$

no matter which finite subset $\widehat{S}_n \subset S_n$ we choose. Suppose now that $A^{-1}a \cap B^{-1}b \cap S_n \neq \emptyset$ and consider the following cases:

(a) $A^{-1}a \cup B^{-1}b \subset S_n$,

(b) $A^{-1}a \cup B^{-1}b \not\subset S_n$.

We will show how to proceed in case (b) (case (a) can be treated in a very similar way). Recall that $S_n = I^{\mathbb{R}}[(2n-1)\widetilde{a}_{n-1}] \times SU(2)$, so in view of the definition of $\mathcal{A}$ one of the following holds:

(b1) $A^{-1}a \cup B^{-1}b \subset (l-2, l+1) \times U_i$,

(b2) $A^{-1}a \cup B^{-1}b \subset (l-1, l+2) \times U_i$,

(b3) $A^{-1}a \cup B^{-1}b \subset (-l-2, -l+1) \times U_i$,

(b4) $A^{-1}a \cup B^{-1}b \subset (-l-1, -l+2) \times U_i$,

[39] The existence of such a sequence can be shown e.g. in the following way. Choose a finite open cover of $\mathbb{T} \times SU(2)$ consisting of continuity sets (i.e. sets whose boundary have zero measure) which are simply connected. It yields a partition of $\mathbb{T} \times SU(2)$ into a finite number of open sets, up to a set of measure zero. Using Theorem 6.5, for each set of this partition, we can find a homeomorphism to $(0,1)^4$ carrying the conditional Haar measure to the Lebesgue measure. Any sequence which is uniformly distributed in $[0,1)^4$ yields sequences uniformly distributed in the closure of the elements of the partition. The desired sequence can be now constructed by taking elements from these sequences with frequencies approximating the measures of the sets from the "cover".

where $1 \le i \le M$ and $l = -(2n-1)\widetilde{a}_{n-1}$. We will cover case (b1) (the other cases can be treated in the same way). We have

$$\begin{aligned}
\lambda_{S_n}(A^{-1}a \cap B^{-1}b) = &\frac{\lambda((l-1,l+2)\times SU(2))}{\lambda(S_n)} \\
&\cdot \lambda_{(l-1,l+2)\times SU(2)}(A^{-1}a \cap B^{-1}b \cap ((l,l+2)\times SU(2))) \\
=&\frac{3}{2|l|}\cdot \lambda_{(l-1,l+2)\times SU(2)}(A^{-1}a \cap B^{-1}b \cap ((l,l+2)\times SU(2))) \\
=&\frac{3}{2|l|}\cdot \lambda\Big(\psi\circ\psi_{l,i}\circ g_a \circ g \circ \psi_{l_A,i_A}^{-1}\circ\psi^{-1}(C_A) \\
&\cap \psi\circ\psi_{l,i}\circ g_b \circ g \circ \psi_{l_B,i_B}^{-1}\circ\psi^{-1}(C_B) \cap \big((1/3,1)\times(0,1)^3\big)\Big) \\
=&\frac{3}{2|l|}\cdot \lambda(\Psi_A(C_A)\cap\Psi_B(C_B)\cap C),
\end{aligned} \tag{91}$$

where

$$\begin{aligned}
\Psi_A &= \psi\circ\psi_{l,i}\circ g_a\circ g\circ\psi_{l_A,i_A}^{-1}\circ\psi^{-1}, \\
\Psi_B &= \psi\circ\psi_{l,i}\circ g_b\circ g\circ\psi_{l_B,i_B}^{-1}\circ\psi^{-1}, \\
C &= (1/3,1)\times(0,1)^3.
\end{aligned}$$

Define $(y_n)_{n\in\mathbb{N}} \subset [0,1]^4$, by

$$y_n = \begin{cases} \psi\circ\psi_{l,i}(x_{(n+2)/3,l-1}), & \text{if } n\equiv 0 \mod 3, \\ \psi\circ\psi_{l,i}(x_{(n+1)/3,l}), & \text{if } n\equiv 1 \mod 3, \\ \psi\circ\psi_{l,i}(x_{n/3,l+1}), & \text{if } n\equiv 2 \mod 3. \end{cases}$$

Notice that $(y_n)_{n\in\mathbb{N}}$ is uniformly distributed in $[0,1]^4$ and the discrepancy of this sequence does not depend on $l\in\mathbb{Z}$. We have

$$\begin{aligned}
&\frac{1}{N}\sum_{n=1}^{N}\mathbb{1}_{A^{-1}a\cap B^{-1}b\cap((l,l+2)\times SU(2))}(x_{n,l-1}) \\
&\qquad+\frac{1}{N}\sum_{n=1}^{N}\mathbb{1}_{A^{-1}a\cap B^{-1}b\cap((l,l+2)\times SU(2))}(x_{n,l}) \\
&\qquad+\frac{1}{N}\sum_{n=1}^{N}\mathbb{1}_{A^{-1}a\cap B^{-1}b\cap((l,l+2)\times SU(2))}(x_{n,l+1}) \\
&=3\cdot\frac{1}{3N}\sum_{n=1}^{3N}\mathbb{1}_{\Psi_A(C_A)\cap\Psi_B(C_B)\cap C}(y_n) \\
&\to 3\lambda(\Psi_A(C_A)\cap\Psi_B(C_B)\cap C).
\end{aligned} \tag{92}$$

In view of Lemma 6.10 the above convergence is uniform with respect to A, B and a, b. It follows from (91) and (92) that $\widehat{S}_n$ defined in the following way:

$$\widehat{S}_n = S_n \cap \left(\bigcup_{l\in Z}\bigcup_{n=1}^{n_0} x_{n,l}\right).$$

satisfies (i) provided that n_0 is large enough. □

Proof of Proposition 6.6 (ii). We claim that the functions from

$$\{f\colon G\times G\to\mathbb{R}\colon f(v,w)=\lambda_{F_n}(Aav\cap Bbw)\}$$
$$\text{are uniformly equi-continuous.}\tag{93}$$

Fix $\varepsilon>0$. Let $\delta\in(0,\delta_0)$ be sufficiently small, so that for $C\in\mathcal{A}$ and $c\in G$ such that $\lambda(Cc)=\lambda(C)>\varepsilon/2$ we can find a ball of radius δ being a subset of Cc. Such choice is possible in view of Lemma 6.10. Take $A,B\in\mathcal{A}$, $a,b\in G$, $n\in\mathbb{N}$ and let $f(v,w)=\lambda_{F_n}(Aav\cap Bbw)$. Take $v_1,v_2,w\in G$ with $d(v_1,v_2)<\delta$. We have

$$|f(v_1,w)-f(v_2,w)|\leq\lambda((Aav_1\triangle Aav_2)\cap Bbw).\tag{94}$$

We will consider the following cases:

(i) $(Aav_1\cup Aav_2)\cap Bbw=\emptyset$,

(ii) $(Aav_1\cup Aav_2)\cap Bbw\neq\emptyset$

  (a) $\lambda(A)\leq\varepsilon/2$,

  (b) $\lambda(A)>\varepsilon/2$.

Using (94) we obtain in case (i) $|f(v_1,w)-f(v_2,w)|=0$ and in case (iia) $|f(v_1,w)-f(v_2,w)|\leq\varepsilon$. In case (iib), by the choice of δ it is clear that $Aav_1\cap Aav_2\neq\emptyset$. Therefore (see the definition of $\widetilde{\mathcal{A}}$) there exist $l\in\mathbb{Z}$ and $1\leq i\leq M$ such that

$$(Aav_1\cup Aav_2\cup Bbw)_{\delta_0}\cup(Aav_1\cup Aav_2\cup Bbw)\subset(l-1,l+2)\times U_i.$$

Now we can use the homeomorphism $\psi_{l,i}^{-1}\circ\psi$ to "transport" the above set to $[0,1]^4$. Using Lemma 6.10 and adjusting δ if necessary we arrive at (93).

Let $K=(2n-1)\widetilde{a}_{n-1}$ and for $k\in[-K,K-1]$ let $y_{m,k}=x_m+(k,I)$. Moreover, for $1\leq i\leq M$ let $(y^i_{m,k})_{m\in\mathbb{N}}$ be a subsequence of elements of $(y_{m,k})_{m\in\mathbb{N}}$ which are in $[k,k+1]\times U_i$ and let $N_i=\#\{x_{m,k}\in U_i\colon 1\leq m\leq N\}$ (notice that this quantity is independent of k). Then for a fixed w we have

$$\begin{aligned}
&\frac{1}{2KN}\sum_{k=-K}^{K-1}\sum_{m=1}^{N}f(y_{m,k},w)\\
&\quad=\frac{1}{2KN}\sum_{k}\sum_{m}\sum_{l}\sum_{i}\eta_l\times\rho_i(y_{m,k})f(y_{m,k},w)\\
&\quad=\frac{1}{2KN}\sum_{i}\sum_{l}\sum_{k=l-1}^{l+1}\sum_{m}\eta_l\times\rho_i(y_{m,k})f(y_{m,k},w)\\
&\quad=\frac{1}{2KN}\sum_{i}\frac{N_i}{N}\sum_{l}\frac{1}{N_i}\sum_{k=l-1}^{l+1}\sum_{m=1}^{N_i}\underbrace{\eta_l\otimes\rho_i(y^i_{m,k})f(y_{m,k},w)}_{f_{l,i,m,k}}.
\end{aligned}\tag{95}$$

We have

$$\frac{1}{3N_i}\sum_{k=l-1}^{l+1}\sum_{m=1}^{N_i}f_{l,i,m,k}(y^i_{m,k})\to\int f_{l,i,m,k}\,d\lambda_{(l-1,l+2)\times SU(2)}$$

and

$$\frac{N_i}{N} \to \lambda_{SU(2)}(U_i)$$

and the above convergences are uniform in view of Theorem 6.3 and Lemma 6.10. Hence the expression (95) converges to

$$\frac{1}{2K}\sum_{i=1}^{M}\sum_{l}\lambda_{SU)(2)}(U_i)\cdot 3\cdot\int f_{l,i,m,k}\ d\lambda_{(l-1,l+2)\times U_i}$$

$$= \frac{1}{2K}\int\sum_{l}\sum_{i=1}^{M}3\lambda_{SU(2)}(U_i)\eta_l\otimes\rho_i\cdot f(\cdot,w)\ d\lambda_{(l-1,l+2)\times U_i}$$

$$= \int f(\cdot,w)\ d\lambda_{S_n}.$$

To end the proof it suffices to use Fubini's theorem and take $\widehat{S}_n$ defined in the following way:

$$\widehat{S}_n = S_n \cap \left(\bigcup_{k\in\mathbb{Z}}\bigcup_{n=1}^{n_0} x_{n,k}\right)$$

for n_0 large enough. Notice that this is consistent with the final choice of $\widehat{S}_n$ in the proof of Proposition 6.6 (i). □

Remark 6.2. Notice that without loss of generality, in Proposition 6.6 we may assume that $F_n \in \mathcal{A}$ for $n \in \mathbb{N}$.

Remark 6.3. The assertion of Proposition 6.6 remains true if instead of sets $A, B \in \mathcal{A}$ we consider sets A', B' which are finite sums of translations of elements from $\mathcal{A}$, provided that the number of elements in the sum is bounded for each n.

References

[1] L. M. Abramov and V. A. Rohlin. Entropy of a skew product of mappings with invariant measure. *Vestnik Leningrad. Univ.*, 17(7):5–13, 1962.

[2] W. Ambrose. Representation of ergodic flows. *Ann. of Math. (2)*, 42:723–739, 1941.

[3] W. Ambrose and S. Kakutani. Structure and continuity of measurable flows. *Duke Math. J.*, 9:25–42, 1942.

[4] T. Austin. Pleasant extensions retaining algebraic structure, i. 05 2009.

[5] J. R. Blum and N. Friedman. On commuting transformations and roots. *Proc. Amer. Math. Soc.*, 17:1370–1374, 1966.

[6] W. Bułatek, M. Lemańczyk, and E. Lesigne. On the filtering problem for stationary random $\mathbb{Z}^2$-fields. *IEEE Trans. Inform. Theory*, 51(10):3586–3593, 2005.

[7] R. V. Chacon. Transformations having continuous spectrum. *J. Math. Mech.*, 16:399–415, 1966.

[8] R. V. Chacon. A geometric construction of measure preserving transformations. In *Proc. Fifth Berkeley Sympos. Math. Statist. and Probability (Berkeley, Calif., 1965/66), Vol. II: Contributions to Probability Theory, Part 2*, pages 335–360. Univ. California Press, Berkeley, Calif., 1967.

[9] R. V. Chacon. Weakly mixing transformations which are not strongly mixing. *Proc. Amer. Math. Soc.*, 22:559–562, 1969.

[10] I. P. Cornfeld, S. V. Fomin, and Y. G. Sinaĭ. *Ergodic theory*, volume 245 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, 1982.

[11] A. I. Danilenko. On simplicity concepts for ergodic actions. *J. Anal. Math.*, 102:77–117, 2007.

[12] A. I. Danilenko. (C, F)-actions in ergodic theory. In *Geometry and dynamics of groups and spaces*, volume 265 of *Progr. Math.*, pages 325–351. Birkhäuser, Basel, 2008.

[13] T. de la Rue. Notes on Austin's multiple ergodic theorem. 07 2009.

[14] T. de la Rue and J. de Sam Lazaro. Une transformation générique peut être insérée dans un flot. *Ann. Inst. H. Poincaré Probab. Statist.*, 39(1):121–134, 2003.

[15] A. del Junco. A simple measure-preserving transformation with trivial centralizer. *Pacific J. Math.*, 79(2):357–362, 1978.

[16] A. del Junco. A simple map with no prime factors. *Israel J. Math.*, 104:301–320, 1998.

[17] A. del Junco and M. Lemańczyk. Simple systems are disjoint from Gaussian systems. *Studia Math.*, 133(3):249–256, 1999.

[18] A. del Junco and M. Lemańczyk. Joinings of distally simple automorphisms. Unpublished preprint, 2005.

[19] A. del Junco and D. Rudolph. On ergodic actions whose self-joinings are graphs. *Ergodic Theory Dynam. Systems*, 7(4):531–557, 1987.

[20] A. del Junco and D. J. Rudolph. A rank-one, rigid, simple, prime map. *Ergodic Theory Dynam. Systems*, 7(2):229–247, 1987.

[21] A. del Junco and R. Yassawi. Multiple mixing and rank one group actions. *Canad. J. Math.*, 52(2):332–347, 2000.

[22] S. Ferenczi. Systèmes de rang un gauche. *Ann. Inst. H. Poincaré Probab. Statist.*, 21(2):177–186, 1985.

[23] S. Ferenczi. Systems of finite rank. *Colloq. Math.*, 73(1):35–65, 1997.

[24] K. Frączek and M. Lemańczyk. On mild mixing of special flows over irrational rotations under piecewise smooth functions. *Ergodic Theory Dynam. Systems*, 26(3):719–738, 2006.

[25] K. Frączek and M. Lemańczyk. Ratner's property and mild mixing for special flows over two-dimensional rotations. *J. Mod. Dyn.*, 4(4):609–635, 2010.

[26] H. Furstenberg. Disjointness in ergodic theory, minimal sets, and a problem in Diophantine approximation. *Math. Systems Theory*, 1:1–49, 1967.

[27] H. Furstenberg. *Recurrence in ergodic theory and combinatorial number theory*. Princeton University Press, Princeton, N.J., 1981. M. B. Porter Lectures.

[28] E. Glasner, B. Host, and D. Rudolph. Simple systems and their higher order self-joinings. *Israel J. Math.*, 78(1):131–142, 1992.

[29] S. Glasner. Quasifactors in ergodic theory. *Israel J. Math.*, 45(2-3):198–208, 1983.

[30] F. Hahn and W. Parry. Some characteristic properties of dynamical systems with quasi-discrete spectra. *Math. Systems Theory*, 2:179–190, 1968.

[31] P. R. Halmos. Square roots of measure preserving transformations. *Amer. J. Math.*, 64:153–166, 1942.

[32] P. R. Halmos. *Lectures on ergodic theory*. Publications of the Mathematical Society of Japan, no. 3. The Mathematical Society of Japan, 1956.

[33] P. R. Halmos. *Lectures on ergodic theory*. Chelsea Publishing Co., New York, 1960.

[34] E. Hlawka. Discrepancy and Riemann integration. In *Studies in Pure Mathematics (Presented to Richard Rado)*, pages 121–129. Academic Press, London, 1971.

[35] A. Kanigowski. Ratner's property for special flows over irrational rotations under functions of bounded variation. 02 2013.

[36] A. Katok. *Combinatorial constructions in ergodic theory and dynamics*, volume 30 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2003.

[37] J. L. F. King. The generic transformation has roots of all orders. *Colloq. Math.*, 84/85(part 2):521–547, 2000. Dedicated to the memory of Anzelm Iwanik.

[38] A. V. Kočergin. The homology of functions over dynamical systems. *Dokl. Akad. Nauk SSSR*, 231(4):795–798, 1976.

[39] U. Krengel and H. Michel. Über Wurzeln ergodischer Transformationen. *Math. Z.*, 96:50–57, 1967.

[40] L. Kuipers and H. Niederreiter. *Uniform distribution of sequences*. Wiley-Interscience [John Wiley & Sons], New York, 1974. Pure and Applied Mathematics.

[41] J. Kułaga-Przymus and F. Parreau. Disjointness properties for Cartesian products of weakly mixing systems. *Colloquium Mathematicum*, 128(2):153–177, 2012.

[42] J. Lamperti. On flows of measure-preserving transformations. *Michigan Math. J.*, 4:161–164, 1957.

[43] M. Lemańczyk and M. K. Mentzen. Compact subgroups in the centralizer of natural factors of an ergodic group extension of a rotation determine all factors. *Ergodic Theory Dynam. Systems*, 10(4):763–776, 1990.

[44] M. Lemańczyk and F. Parreau. Rokhlin extensions and lifting disjointness. *Ergodic Theory Dynam. Systems*, 23(5):1525–1550, 2003.

[45] M. Lemańczyk, F. Parreau, and E. Roy. Joining primeness and disjointness from infinitely divisible systems. *Proc. Amer. Math. Soc.*, 139(1):185–199, 2011.

[46] M. Lemańczyk, F. Parreau, and J.-P. Thouvenot. Gaussian automorphisms whose ergodic self-joinings are Gaussian. *Fund. Math.*, 164(3):253–293, 2000.

[47] M. Lemańczyk, J.-P. Thouvenot, and B. Weiss. Relative discrete spectrum and joinings. *Monatsh. Math.*, 137(1):57–75, 2002.

[48] E. Lindenstrauss. Pointwise theorems for amenable groups. *Invent. Math.*, 146(2):259–295, 2001.

[49] M. K. Mentzen. Ergodic properties of group extensions of dynamical systems with discrete spectra. *Studia Math.*, 101(1):19–31, 1991.

[50] H. Michel. Wurzeln vollergodischer Automorphismen mit quasidiskretem Spektrum. I. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 9:323–340, 1968.

[51] H. Michel. Vollergodische Automorphismen mit quasidiskretem Spektrum und Wurzeln jeder rationalen Ordnung. *Wiss. Beiträge Martin-Luther-Univ. Halle-Wittenberg*, M1:123–128, 1968/1969.

[52] H. Michel. Wurzeln vollergodischer Automorphismen mit quasidiskretem Spektrum. II. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 16:321–335, 1970.

[53] J. Neveu. *Processus aléatoires gaussiens*. Séminaire de Mathématiques Supérieures, No. 34 (Été, 1968). Les Presses de l'Université de Montréal, Montreal, Que., 1968.

[54] D. Newton. On canonical factors of ergodic dynamical systems. *J. London Math. Soc. (2)*, 19(1):129–136, 1979.

[55] D. Ornstein. Bernoulli shifts with the same entropy are isomorphic. *Advances in Math.*, 4:337–352 (1970), 1970.

[56] D. Ornstein. Two Bernoulli shifts with infinite entropy are isomorphic. *Advances in Math.*, 5:339–348 (1970), 1970.

[57] D. S. Ornstein. Imbedding Bernoulli shifts in flows. In *Contributions to Ergodic Theory and Probability (Proc. Conf., Ohio State Univ., Columbus, Ohio, 1970)*, pages 178–218. Springer, Berlin, 1970.

[58] D. S. Ornstein. On the root problem in ergodic theory. In *Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability (Univ. California, Berkeley, Calif., 1970/1971), Vol. II: Probability theory*, pages 347–356, Berkeley, Calif., 1972. Univ. California Press.

[59] D. S. Ornstein. The isomorphism theorem for Bernoulli flows. *Advances in Math.*, 10:124–142, 1973.

[60] D. S. Ornstein. A K automorphism with no square root and Pinsker's conjecture. *Advances in Math.*, 10:89–102, 1973.

[61] M. Osikawa. Centralizer of an ergodic measure preserving transformation. *Publ. Res. Inst. Math. Sci.*, 18(1):35–48, 1982.

[62] J. C. Oxtoby and S. M. Ulam. Measure-preserving homeomorphisms and metrical transitivity. *Ann. of Math. (2)*, 42:874–920, 1941.

[63] W. Parry. Zero entropy of distal and related transformations. In *Topological Dynamics (Symposium, Colorado State Univ., Ft. Collins, Colo., 1967)*, pages 383–389. Benjamin, New York, 1968.

[64] M. Ratner. Factors of horocycle flows. *Ergodic Theory Dynam. Systems*, 2(3-4):465–489 (1983), 1982.

[65] M. Ratner. Rigidity of horocycle flows. *Ann. of Math. (2)*, 115(3):597–614, 1982.

[66] M. Ratner. Horocycle flows, joinings and rigidity of products. *Ann. of Math. (2)*, 118(2):277–313, 1983.

[67] D. J. Rudolph. An example of a measure preserving map with minimal self-joinings, and applications. *J. Analyse Math.*, 35:97–122, 1979.

[68] V. V. Ryzhikov and J.-P. Thouvenot. Disjointness, divisibility, and quasi-simplicity of measure-preserving actions. *Funktsional. Anal. i Prilozhen.*, 40(3):85–89, 2006.

[69] V. V. Ryzhikov and S. V. Tikhonov. Generic $\mathbb{Z}^n$-action can be embedded only in injective $\mathbb{R}^n$-actions. *Mat. Zametki*, 79(6):925–930, 2006.

[70] A. Siemaszko. On group extensions of 2-fold simple ergodic actions. *Studia Math.*, 110(1):53–64, 1994.

[71] A. M. Stepin and A. M. Eremenko. Nonuniqueness of an inclusion in a flow and the vastness of a centralizer for a generic measure-preserving transformation. *Mat. Sb.*, 195(12):95–108, 2004.

[72] J.-P. Thouvenot. Some properties and applications of joinings in ergodic theory. In *Ergodic theory and its connections with harmonic analysis (Alexandria, 1993)*, volume 205 of *London Math. Soc. Lecture Note Ser.*, pages 207–235. Cambridge Univ. Press, Cambridge, 1995.

[73] S. V. Tikhonov. The generic action of the group $\mathbb{Z}^d$ can be embedded into the action of the group $\mathbb{R}^d$. *Dokl. Akad. Nauk*, 391(1):26–28, 2003.

[74] S. V. Tikhonov. Embeddings of lattice actions in flows with multidimensional time. *Mat. Sb.*, 197(1):97–132, 2006.

[75] W. A. Veech. A criterion for a process to be prime. *Monatsh. Math.*, 94(4):335–341, 1982.

[76] D. Witte. Rigidity of some translations on homogeneous spaces. *Invent. Math.*, 81(1):1–27, 1985.

[77] R. J. Zimmer. Ergodic actions with generalized discrete spectrum. *Illinois J. Math.*, 20(4):555–588, 1976.

[78] R. J. Zimmer. Extensions of ergodic group actions. *Illinois J. Math.*, 20(3):373–409, 1976.

[79] R. J. Zimmer. Induced and amenable ergodic actions of Lie groups. *Ann. Sci. École Norm. Sup. (4)*, 11(3):407–428, 1978.

[80] R. J. Zimmer. *Ergodic theory and semisimple groups*, volume 81 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1984.

Joanna Kułaga-Przymus
Faculty of Mathematics and Computer Science
Nicolaus Copernicus University
ul. Chopina 12/18, 87–100 Toruń, Poland
joanna.kulaga@gmail.com